

# Gamma-X Event detection And Work at SLAC

Jonathan Nikoleyczik


I am attempting to understand a potential background in the LZ detector, called Gamma-X events by analyzing simulations of background events.

Today's update starts on [slide 15](#)

# Running BACCARAT

- Ran test Sims in latest version of BACCARAT (1.0.0)
- Will modify scripts to look for Gamma-X events
- Currently using simple rootpy analysis
- Plan to emulate Bhawna's Gamma-X study


S1 vs. S2 from the BACCARAT test macro



# More Gamma-X results


- 100,000 Th 232 Decay Chain Events
- Expected an exponential drop off ( I see one if I don't require energy deposited)
- 

Events in both Forward and Reverse Field Regions



# Gamma-X Update


- Same 100,000 Events as last week
- Producing 560 events in both regions
- Now I add together events in forward and reverse field regions, then do the energy weighted sum
- There is now one dot for each decay event
- Issues getting S1 and S2
  - Needs libNEST
  - Needs Boost



# Fixed Histogram

- This is a histogram of the data on the last slide as a function of  $z$
- The RFR looks almost linear and the FFR looks exponential


Energy weighted events in both forward and reverse field regions  
For individual primary particles



# fastNEST is up and running


I'm able to get S1 and S2s

The plot on the right is just an example of S1 and S2 for Nuclear and Electronic events. They are for random energy depositions of 1 to 100 keV with no specification on location.




# Real Gamma-X events in S1 and S2

- Gamma events create a lot of light compared to WIMP events
- See up to 6000 S1 but the WIMP search is only up to 50 S1
- See the expected suppression of S2



# Just WIMP search region


- Same data, now only showing the WIMP search region defined in the TDR
- See ~5 events that appear to be closer to nuclear rather than electronic events.
- Need more statistics, this is with 100,000 decay events I can easily scale this up to a few million events





# Search Region from TDR

This is the plot which I used to compare the normal and Gamma-X events to.




**Figure 1.3.11:** Discrimination parameter  $\log_{10}(S2/S1)$  as a function of S1 signal obtained with LUX calibration [58]. (a) ER band calibrated with beta decays from a dispersed  ${}^3\text{H}$  source; the median is shown in blue, with 80% population contours indicated by the dashed blue lines. (b) NR band populated by elastic neutron scattering from a D-D pulsed neutron source; the median (solid) and 80% band width (dotted) are indicated in blue and red, respectively. The mostly vertical gray lines are contours of constant energy deposition. For more information, see Chapter 7.

# Time resolution

I wanted to make sure that all of the events occur within the timeframe of one event in the detector


Almost all events occur within a few nanoseconds of the initial decay.

There is a much smaller population of secondary events which are handled separately from the primary events.



# High statistics run


- Went from 100,000 decays to 10,000,000 decays in this data set
- There are band like structures in the  $S1$  vs  $\log(S2/S1)$  plot (shown right)
- I am only focusing on the WIMP search region
- There are ~12,000 Gamma-X events in this region



# Including ER and NR bands

The Gamma-X events do not exactly follow the ER and NR bands. There is suppression of S2 at constant S1.


The bands shown here are the 90% bands.



# Focusing just on the NR band

There are 1098 events that fall in this band.

So about 0.01% of Th232 decays from the PMT windows will be likely to cause Gamma-X events



# Next Steps

- More high statistics runs of other radioactive sources. Likely Uranium as it also has a high rate from the PMT windows.
- Detailed analysis of these events. Likely just to determine the tracks and angles which create these events (Are they more likely in the edge or the center etc.).
- Determine the actual rate in cts/kg/day. This is likely to be an underestimation as there are more background sources than the PMT windows

# Gamma-X Events

Jonathan Nikoleyczik  
UW - Madison  
Physics Meeting


# Overview

- Simulate 20,000,000 Th232 decay chain events created at the PMT windows
  - Used BACCARAT 1.2.0 with BaccRootConverter
- Calculate S1 and S2 using libNEST
- Analysis code to look for events which deposit energy in both the forward and reverse field regions (Gamma-X events)
  - [Jupyter Notebook Analysis Code](#) (First method)
  - [Analysis Code](#) (Corrected method)
  - [BACCARAT macro](#)




# Gamma-X events

A diagram of a Gamma-X event. An interaction occurs in the reverse field region which creates an S1 but no corresponding S2. The track continues into the forward field region and creates a normal S1 and S2 which will have a suppressed S2 compared to the total energy deposited.




# Analysis method

- Follow the method of the TDR ReducedAnalysisTree
- Group together events within a small region then calculate S1 and S2 based on the total energy deposited in that region
- Moves events out of the WIMP search region significantly reducing the rates
- Also decreases the S2 to S1 ratio



# WIMP Search Region

- Since S1 and S2 are calculated with higher energy the S1 and S2 move out to higher values and we get fewer events in the search region



# Improved method rates

| Source | Decays Simulated | Fraction which produce Gamma-X (all) | Fraction in WIMP search region (Depending on S1 S2 cut) | Approximate rate (assuming production from PMT windows and <b>no cuts</b> ) |
|--------|------------------|--------------------------------------|---|---|
| Th-232 | 14,900,000       | 0.012887<br>192507 Events            | $\sim 4.0 \times 10^{-7}$                               | 0.24 events per year  |
| U-238  | 4,150,000        | 0.013739<br>57094 Events             | $\sim 1.2 \times 10^{-6}$                               | 3.83 events per year  |
| Co-60  | 9,800,000        | 0.080683<br>790704 Events            | $\sim 4.0 \times 10^{-7}$                               | 0 (No Co60 in PMT windows)  |
| K-40   | 9,400,000        | 0.004208<br>39559 Events             | $< 1.0 \times 10^{-7}$                                  | $< 0.13$ events per year  |

Agrees with rates from previous simulations of Gamma-X rates [Bhwana Gomers slides](#)


# Future

- Add cuts
  - Follow TDR cuts (energy, time, fiducial volume)
  - Preliminary but fiducial volume gets rid of almost all events
  - See if there are more that would be useful
- Look at decays from more sources
  - Include walls and PMT holder


Backup Slides  
Continued on Slide 36:  
Work at SLAC

# Energy deposits agree with Th 232 spectrum


Thorium Spectrum (2"x2" Ludlum 44-10)



Gamma-X Energy Deposited Th232 No Cuts




# Normal Events (Th 232)





# The First Method


- Focus on only the Gamma-X events
- For each energy deposition calculate an S1 and S2. Then sum S1 and S2 for each event.
- This caused this band-like structure as energy depositions have sharp lines at very low energy ( $\sim 1$  keV).



# Energy Depositions

There are very sharp lines in energy deposited which are more common in the banded S1/S2 events.



Energies correspond to Xe Energy Levels 4.78, 5.10, and 5.45 keV



# Approximate rate in first method


- Simulated 20,000,000 Decay chain events from the bottom PMTs
- 12,000 (0.6%) within the DM search region (0-50 S1 and 1.2-3.8 in  $\log(S2/S1)$ )
- 2,900 events (0.15%) fall below the 90% ER band (blue dotted line on previous plot)
- 1,100 events (0.05%) are within the NR band
- This corresponds to 2.5 events per day. Note that this is before any cuts are made. The fiducial volume cut will reduce this number significantly. And this is only from events originating in the bottom PMTs
- This is still much much larger than we expected from previous background studies

# Th 232 Results (Improved Method)




# Th 232 Results



Gamma-X Energy Deposited Th232 No Cuts



Gamma-X Events Doke Plot Th232 No Cuts




# U 238 Results




# U 238 Results



Gamma-X Energy Deposited U 238 No Cuts



Gamma-X Events Doke Plot U 238 No Cuts




# Co 60 Results






# Co 60 Results

Gamma-X Energy Deposited Co60 No Cuts




Gamma-X Events Doke Plot Co60 No Cuts




# K 40 Results

Gamma-X Events K40 No Cuts




Gamma-X Events K 40 No Cuts




# K 40 Results

Gamma-X Energy Deposited K 40 No Cuts




Gamma-X Events Doke Plot K 40 No Cuts



# Work at SLAC

- Almost finished the gas test hood
- Cut pieces of T-slot and plexiglass for the phase 2 clean room
- Disassembled the phase 1 breakouts for reassembly
  - We are redesigning the breakout to accommodate new thermosyphon lines which will be installed next week



Gas test hood drawing