



Multigap Diffraction at the LHC

K. Goulianos

The Rockefeller University




DIS 2005

27 April - 1 May
Madison, Wisconsin

<http://physics.rockefeller.edu/dino/my.html>


Multigap Diffraction

(KG, hep-ph/0205141)



Elastic and Total Cross Sections


QCD expectations



$$\sigma_T(s) = \sigma_0 s^\varepsilon = \sigma_0 e^{\varepsilon \Delta y'}$$


The exponential rise of $\sigma_T(\Delta y')$ is due to the increase of wee partons with $\Delta y'$

(see E. Levin, An Introduction to Pomeron, Preprint DESY 98-120)





$$\text{Im } f_{el}(s, t) \propto e^{\varepsilon + \alpha' t \Delta y}$$

Total cross section:
power law rise with energy



Elastic cross section:
forward scattering amplitude

Single Diffraction



2 independent variables: $t, \Delta y$

$$\frac{d^2\sigma}{dt d\Delta y} = \underbrace{C \cdot F_p^2(t) \cdot e^{|\varepsilon + \alpha' t| \Delta y}}_{\text{gap probability}} \cdot \underbrace{\mathbf{K} \cdot \left| \sigma_o e^{\varepsilon \Delta y'} \right|}_{\text{sub-energy x-section}}$$

color
factor

$$\mathbf{K} = \frac{g_{IP-IP-IP}(t)}{\beta_{IP-p-p}(0)} \approx 0.17$$

Gap probability MUST be normalized to unity!

Single diffraction (re)normalized

$$\frac{d^2\sigma}{dt d\Delta y} = N_{gap} \cdot \underbrace{C \cdot F_p^2(t) \cdot e^{|\varepsilon + \alpha' t| \Delta y}}_{P_{gap}(\Delta y, t)} \cdot \kappa \cdot \sigma_o e^{\varepsilon \Delta y'}$$


$$N_{gap}^{-1}(s) = \int_{\Delta y, t} P_{gap}(\Delta y, t) d\Delta y dt \xrightarrow{s \rightarrow \infty} C' \cdot \frac{s^{2\varepsilon}}{\ln s}$$

$$\frac{d^2\sigma}{dt d\Delta y} = C'' \left| e^{\varepsilon(\Delta y - \ln s)} \cdot \ln s \right| e^{(b_0 + 2\alpha' \Delta y)t}$$

Grows slower than s^ε

→ The Pumplin bound is obeyed at all impact parameters

Total Single Diffractive x-Section



$$\frac{d^2\sigma_{SD}}{dtd\xi} = f_{IP/p}(t, \xi) \cdot \sigma_{IP-\bar{p}}(s\xi)$$

$$\sigma_{SD} \sim s^{2\epsilon}$$

❖ Unitarity problem:


Using factorization and std pomeron flux σ_{SD} exceeds σ_T at $\sqrt{s} \approx 2$ TeV.

❖ Renormalization:

Normalize the Pomeron flux to unity

KG, PLB 358 (1995) 379

$$\int_{\xi_{min}}^{0.1} \int_0^{\infty} f_{IP/p}(t, \xi) d\xi dt = 1$$



The Factors κ and ε



Experimentally:

KG&JM, PRD 59 (114017) 1999

$$\kappa = \frac{g_{IP-IP-IP}}{\beta_{IP-p}} = 0.17 \pm 0.02, \quad \varepsilon = 0.104$$

Color factor: $\kappa = f_g \times \frac{1}{N_c^2 - 1} + f_q \times \frac{1}{N_c} \frac{Q^2 = 1}{Q^2 = 1} \rightarrow \approx 0.75 \times \frac{1}{8} + 0.25 \times \frac{1}{3} = 0.18$

Pomeron intercept: $\varepsilon = \lambda_g \cdot W_g + \lambda_q \cdot W_q = 0.12$




$$X \cdot f(X) = \frac{1}{X^\lambda}$$

f_g = gluon fraction
 f_q = quark fraction


$$\int_{x=1/s}^1 f(x) dx \sim s^\lambda$$

Multigap Cross Sections



5 independent variables

$$\left\{ \begin{array}{l} t_1 \\ \Delta y = \Delta y_1 + \Delta y_2 \end{array} \right.$$




color factors

$$\prod_{i=1-5} \frac{d^5 \sigma}{dV_i} = C \times F_p^2(t_1) \prod_{i=1-2} \left| e^{|\varepsilon + \alpha' t_i| \Delta y_i} \right|^2 \times \underbrace{\kappa^2 \sigma_o e^{\varepsilon |\Delta y'_1 + \Delta y'_2|}}_{\text{Sub-energy cross section (for regions with particles)}}$$

Gap probability


$$\int_{\Delta y, t} \sim s^{2\varepsilon} / \ln s$$

Same suppression as for single gap!

 DP

➤ Two-gap/one-gap ratios are $\approx \kappa = 0.17$

Gap Survival Probability



$$S = \frac{\phi \eta \phi}{\phi \eta \phi}$$

$$S_{2\text{-gap}/1\text{-gap}}^{1\text{-gap}/0\text{-gap}} (1800 \text{ GeV}) \approx 0.23$$


$$S_{2\text{-gap}/1\text{-gap}}^{1\text{-gap}/0\text{-gap}} (630 \text{ GeV}) \approx 0.29$$

[Results similar to predictions by:](#)
[Gotsman-Levin-Maor](#)
[Kaidalov-Khoze-Martin-Ryskin](#)
[Soft color interactions](#)


Diffractive DIS @ HERA

Factorization holds: J. Collins

Pomeron exchange




Color reorganization



$$F_2^{D(3)}(\xi, x, Q^2) \propto \frac{1}{\xi^{1+\epsilon}} \cdot F_2(x, Q^2)$$


Inclusive vs Diffractive DIS

KG, “Diffraction: a New Approach,” J.Phys.G26:716-720,2000 e-Print Archive: hep-ph/0001092



$$F_2^{D(3)}(\xi, \beta, Q^2) \propto \frac{1}{\xi^{1+\epsilon}} \cdot \frac{C(Q^2)}{(\beta\xi)^\lambda \lambda(Q^2)} \propto \frac{1}{\xi^{1+\epsilon+\lambda}} \cdot \frac{C}{\beta^\lambda}$$

Diffractive Dijets @ Tevatron



$$F^D(\xi, x, Q^2) \propto \frac{1}{\xi^{1+2\varepsilon}} \cdot F(x/\xi, Q^2)$$

$F^D_{JJ}(\xi, \beta, Q^2) @ \text{Tevatron}$

$$F^D(\xi, \beta, Q^2) = N_{\text{renorm}} \frac{1}{\xi^{1+2\varepsilon}} \cdot \frac{C(Q^2)}{(x/\xi)\lambda(Q^2)} = \left(\frac{2\varepsilon}{\beta s} \right)^{2\varepsilon} \cdot \frac{1}{\xi^{1+2\varepsilon}} \cdot \frac{C(Q^2)}{\beta \lambda(Q^2)}$$

$$N_{\text{renorm}}^{-1} = \int_{\xi_{\min}}^1 \frac{d\xi}{\xi^{1+2\varepsilon}}$$

$$\xi_{\min} = \frac{x_{\min}}{\beta} \approx \frac{1}{\beta s}$$

$$\left(\frac{\beta s}{2\varepsilon} \right)^{2\varepsilon}$$


$$\text{RENORM} \Rightarrow R \frac{SD}{ND} \left| x \right| = \frac{2\varepsilon}{s^{2\varepsilon}} \frac{1}{\xi^{1-\lambda(Q^2)}} \cdot x^{-(2\varepsilon)}$$

$$\varepsilon_g = 0.2 \rightarrow x^{-0.4}$$

SD/ND Dijet Ratio vs x_{Bj} @ CDF



$$R(x) = \frac{F_{jj}^{SD}(x)}{F_{jj}^{ND}(x)}$$





$$0.035 < \xi < 0.095$$

Flat ξ dependence

$$R(x) = x^{-0.45}$$




Storing Factorization @ Tevatron




$R(SD/ND)$


$R(DPE/SD)$

DSF from two/one gap:
factorization restored!







Low-x and Diffraction



Derive diffractive
from inclusive PDFs
and color factors



$$x \cdot f(x) = \frac{1}{\epsilon \text{ (or } \lambda \text{)}}$$



Summary

- Multigap processes offer the opportunity of studying diffraction without complications arising from screening corrections, multi-Pomeron exchanges, rescattering, or other rapidity gap survival issues.
- Run 1 results from the Tevatron, and those expected from Run 2, should be followed up by more studies at the LHC with the aim of advancing a successful phenomenology to a THEORY of diffraction.



Thanks to The Rockefeller HEP Group



Anwar



Luc



Stefano



Michele




Mary



Koji



Christina



Andrea



Ken

