Vector meson electroproduction at small Bjorken-x and GPDs

P. Kroll

Fachbereich Physik, Universität Wuppertal

Madison, April 2005

Outline:

- **Handbag factorization and GPDs**
- **Structure of the amplitudes**
- **Modelling the GPDs**
- **The partonic subprocess**
- **Comparison with experiment**
- **Summary**

based on work done in collaboration with

S.V. Goloskokov, hep-ph/0501242

Handbag factorization

Radyushkin (96); Collins et al (97):

proof of factorization

for $Q^2 \to \infty$ into

$\gamma^* g \to V g \;\; (+ \gamma^* q \to V q)$

and GPDs

(at small $x_{\rm Bj}$)

$x \neq x'$



dominant transition $\gamma^*_L \to V_L$ (others power suppressed)

lead. $\ln(1/x_{\rm Bj})$ appr. (Brodsky et al): $x \simeq x' \simeq x_{\rm Bj}$; GPD $\to$ PDF

Generalized Parton Distributions

D. Müller et al (94), Ji(97), Radyushkin (97)

$$\xi = \frac{(p-p')^+}{(p+p')^+} \qquad \bar{x} = \frac{\overline{k}^+}{\overline{p}^+}$$

$$x = \frac{\bar{x}+\xi}{1+\xi} \qquad x' = \frac{\bar{x}-\xi}{1-\xi}$$



$$\int \frac{dz^-}{\pi}\, e^{i\,\bar{x}\,\overline{p}^+ z^-}\, \langle p'|\, G^{+\mu}(-\overline{z}/2)\, G^+_\mu(\overline{z}/2)\, |p\rangle =$$

$$\overline{u}(p')\gamma^+ u(p) H^g(\bar{x},\xi;t) + \overline{u}(p') i\sigma^{+\alpha} \frac{\Delta_\alpha}{2m} u(p)\; E^g(\bar{x},\xi;t)$$

(gauge $A^+ = 0$; $\overline{z} = [0, z^-, \mathbf{0}_\perp]$) $\qquad \widetilde{G}^+_\mu \longrightarrow \widetilde{H}^g, \widetilde{E}^g$

reduction formulas

$H^g(\bar{x},0;0) = \bar{x} g(\bar{x}) \qquad \widetilde{H}^g(\bar{x},0;0) = \bar{x}\Delta g(\bar{x})$

universality, polynomiality, evolution, positivity constraints, Ji's sum rule

The $\gamma^* p \to V p$ amplitudes

kinematical regime: large Q^2, W, small $x_{\rm Bj} (\lesssim 10^{-2})$, t;
kinematics fixes skewness: $\xi \simeq x_{\rm Bj}/2$

$$M_{\mu' +,\mu +} = \frac{e}{2} c_V \int_0^1 \frac{d\bar{x}}{(\bar{x}+\xi)(\bar{x}-\xi+i\epsilon)} \times$$
$$\left\{ \sum_\lambda H^V_{\mu'\lambda,\mu\lambda} [H^g - \frac{\xi^2}{1-\xi^2} E^g] + \sum_\lambda \lambda H^V_{\mu'\lambda,\mu\lambda} [\widetilde{H}^g - \frac{\xi^2}{1-\xi^2} \widetilde{E}^g] \right\}$$

$$M_{\mu' -,\mu +} = \cdots \frac{\sqrt{-t}}{2m} \quad \cdots E^g + \cdots \xi \widetilde{E}^g$$

Unpolarized protons: no flip-nonflip interference

$$|M_{\mu' -,\mu +}|^2 \propto t/m^2 \qquad \text{neglected}$$

$M = M^H + M^{\widetilde{H}}$: $\quad M^{H(\widetilde{H})}_{++,++} = +(-) M^{H(\widetilde{H})}_{-+,-+}$

$$M^{H(\widetilde{H})}_{0+,++} = -(+) M^{H(\widetilde{H})}_{0+,-+}$$

as for (un)natural parity exchange

$\Rightarrow Obs \sim |M^H|^2 + |M^{\widetilde{H}}|^2 \qquad \frac{\widetilde{H}}{H} \simeq \frac{|\Delta g|}{g} \ll 1$

Electroprod. with unpolarized protons at small $x_{\rm Bj}$ and small t probes H^g

Modelling the GPD

GPD controlled by non-pert. QCD: lattice QCD (not yet)
extraction from experiment (not yet)
modelling

Factorizing in $\bar{x}, \xi$ and t probably incorrect (lattice, overlap)
restriction to $t \simeq 0$
Popular ansatz (Radyushkin) $n = 1, 2$

$$f(\beta, \alpha, t \simeq 0) = g(\beta) \frac{\Gamma(2n+2)}{2^{2n+1}\Gamma^2(n+1)} \frac{[(1-|\beta|)^2 - \alpha^2]^n}{(1-|\beta|)^{2n+1}}$$

double distribution; guarantees polynomiality of moments in ξ

$$H^g(\bar{x}, \xi) = \left[\Theta(0 \leq \bar{x} \leq \xi) \int_{\frac{\bar{x}-\xi}{1+\xi}}^{\frac{\bar{x}+\xi}{1+\xi}} + \Theta(\xi \leq \bar{x} \leq 1) \int_{\frac{\bar{x}-\xi}{1-\xi}}^{\frac{\bar{x}+\xi}{1+\xi}} \right] d\beta \frac{\beta}{\xi} f(\beta, \alpha = \frac{\bar{x}-\beta}{\xi})$$

$$+ \xi D^g(\bar{x}/\xi) \qquad [\text{support} - \xi \leq \bar{x} \leq \xi]$$

Input: NLO CTEQ5M



$n = 1$ ($n = 2$ similar)

solid(dashed, dash-dotted) line: $\xi = 5(1, 0.5) \cdot 10^{-3}$

The subprocess $\gamma^* g \to q\bar{q}g$, $q\bar{q} \to V$

modified perturbative approach (Sterman et al):
quark transverse momenta and gluonic radiative corrections (Sudakov) are taken into account;
supresses end-point regions ($\tau \to 0, 1$) where q and $\bar{q}$ are separated by large transerse distances



similar to treatment in lead. $\ln(1/x_{\rm Bj})$ appr. (e.g. Frankfurt et al)

$$H_i^V = \int d\tau \frac{d^2k_\perp}{16\pi^2}\Psi_{Vi}(\tau,k_\perp) \times \frac{1}{\sqrt{2}} Tr\left[(\not{q}' + m_V)\not{\epsilon}_V T_0 - \frac{k_\perp^2 g_\perp^{\alpha\beta}}{2M_V}\{(\not{q}' + m_V)\not{\epsilon}_V, \gamma_\alpha\}\Delta T_\beta + \ldots\right]$$

M_V soft parameter of order m_V
Gaussian wavefunction: $\Psi_{Vi} \sim f_{Vi} \exp\left[-a_{Vi}^2 k_\perp^2/(\tau(1-\tau)\right]$
parameters e.g. $f_{\rho L} = 0.216$ GeV $\quad a_{\rho_L} = 0.52\,\mathrm{GeV}^{-1}$

Hierarchy:

$(\langle k_\perp^2 \rangle^{1/2}/m_V \sim O(1))$

$$
\begin{aligned}
\mathrm{L} \to \mathrm{L} \qquad & H^V \propto 1 \\
\mathrm{T} \to \mathrm{T} \qquad & \propto \langle k_\perp^2 \rangle^{1/2}/Q \\
\mathrm{T} \to \mathrm{L} \qquad & \propto \sqrt{-t}/Q \\
\mathrm{L} \to \mathrm{T} \qquad & \propto \sqrt{-t}\langle k_\perp^2 \rangle^{1/2}/Q^2 \\
\mathrm{T} \to -\mathrm{T} \qquad & \propto -t\langle k_\perp^2 \rangle^{1/2}/Q^3
\end{aligned}
$$

t dependence

Integrated cross sections (σ_L, σ_T)

exponential slopes of amplitudes $\sim e^{t\, B^V_{LL,TT}/2}$

i.e. we essentially need forward amplitudes (calculated)

and slopes (taken from experiment)

$B^V_{TT} \neq B^V_{LL}$? $\qquad |M_{TT}|^2 \propto \left(\frac{f_T^V}{M_V}\right) \frac{1}{B^V_{TT}}$

without precise t dependent data ($d\sigma/dt$, SDME) at disposal

only this product of parameters is probed

Examples:

$B^V_{TT} \simeq B^V_{LL}/2 \qquad M_V = m_V \qquad f_T^\rho = 250$ MeV

$B^V_{TT} \simeq B^V_{LL} \qquad M_V = m_V/2 \qquad f_T^\rho = 170$ MeV

First example slightly favored by SDME

Improvements are required: concept clear but we need $H^g(\bar{x}, \xi, t)$



$W \simeq 75\,\mathrm{GeV}$; filled (open) symbols: H1 (ZEUS) data

solid (dashed) line: ratio of differential (integrated) cross sections, $\tilde{R}\,(R)$

($t = -0.15\,\mathrm{GeV}^2$; $B^V_{TT} = B^V_{LL}/2$)

Spin density matrix elements of the ρ



$W \simeq 75\,\mathrm{GeV}$ and $t \simeq -0.15\,\mathrm{GeV}^2$; filled (open) symbols: H1 (ZEUS) data

$$\sigma_L \sim \frac{|H^g(\xi,\xi)|^2}{B^V_{LL} Q^6} \sim W^{4\delta(Q^2)}$$

for fixed Q^2

from $\bar{x} g(\bar{x}) \sim \bar{x}^{-\delta}$ for $\bar{x} \to 0$



lead. $\ln(1/x_{\mathrm{Bj}})$ approx.:

$H^g(\xi,\xi) \to 2\xi g(2\xi)$

reasonable approx. at small ξ

ca. 18% enhancement

skewing effect (Martin et al, Belitsky et al)



Summary

- We analyzed electroproduction of light vector mesons at large $Q^2 (\gtrsim 4\,\mathrm{GeV}^2)$ and small $x_{\mathrm{Bj}} (\lesssim 10^{-2})$ within the handbag approach
- Only the gluonic GPD H^g contributes
- Subprocess is calculated within modified perturbative approach
- Fair agreement with integrated cross section and SDME is obtained with a few free parameters (essentially for the $T \to T$ transition)
- Amplitudes are calculated at $t = 0$, t dependence assumed to be exponential
- Accurate t dependent data ($d\sigma/dt$, SDME) required for improvement