



Status Report of NNLO QCD Calculations

Michael Klasen
LPSC Grenoble
April 28, 2005



Structure Functions (1)

✓ Motivation:

- Match experimental precision of final HERA data
- Reduce PDF uncertainties for new physics searches at LHC

✓ Complete calculation of $F_{1,2,3}(x, Q^2)$ and $F_L = F_2 - 2xF_1$:

- Anomalous dimensions: Σ @ 3-loop [hep-ph/0404111]
→ Coefficients of $1/\varepsilon$ -poles in dim. reg.
- Coefficient functions: c_l @ 3-loop [hep-ph/0411112]
→ Coefficients of ε^0 -terms in dim. reg.

✓ Very large effort:

- 20 man-years: S. Moch, J. Vermaseren, A. Vogt (1997-2004)
- Advances in mathematics ($H_{1..i}$) and computer algebra (FORM)
- Applications to photon structure functions, e^+e^- -annihilation

Structure Functions (2)

Methods:

- Optical theorem:

$$W_{\mu\nu} = e_{\mu\nu} \frac{1}{2x} F_L(x, Q^2) + d_{\mu\nu} \frac{1}{2x} F_2(x, Q^2) + i \varepsilon_{\mu\nu\alpha\beta} \frac{p^\alpha q^\beta}{p \cdot q} F_3(x, Q^2)$$

- Mellin space:

$$F_i^{\gamma, N}(Q^2) = \int_0^1 dx x^{N-1} \mathcal{F}_i^\gamma(x, Q^2), \quad \mathcal{F}_i(x) = \frac{1}{x} F_i(x)$$


$$F_i^{\gamma, N}(Q^2) = \sum_{\alpha=ns, q, g, \gamma} C_{i, \alpha}^N \left(\frac{Q^2}{\mu^2}, a_s, a_{em} \right) A_{\Gamma, \alpha}^N(\mu^2)$$

Topology mapping:

- Composite → Basic building blocks
- Non-planar → Benz → Ladder

Integral relations:

- Integration by parts
- Scaling identities
- Form factors relations



$$\gamma = \begin{pmatrix} \gamma_{ns} & 0 & 0 & k_{ns} \\ 0 & \gamma_{qq} & \gamma_{qg} & k_{qg} \\ 0 & \gamma_{gq} & \gamma_{gg} & k_{gg} \\ 0 & 0 & 0 & 0 \end{pmatrix}$$




$$\int \prod_n d^D l_n \frac{\partial}{\partial l_i^\mu} l_j^\mu \times (...) = 0$$

$$Q^\mu \frac{\partial}{\partial Q^\mu}, \quad P^\mu \frac{\partial}{\partial Q^\mu}, \quad P^\mu \frac{\partial}{\partial P^\mu}$$


$$\int \prod_n d^D l_n l_i^\mu \times (...) = Q^\mu I_Q + P^\mu I_P$$

Structure Functions (3)

Splitting functions:





Coefficient functions:



Structure Functions (4)

MRST2002: Approx. NNLO

MRST2004: Exact NNLO



Jets (1)

✓ Motivation:

- Determination of α_s and PDFs at LEP, HERA and Tevatron
- Reduce background uncertainties for jet signatures at LHC

✓ So far $2 \rightarrow 2$ processes only:

- $e^+e^- \rightarrow 3 \text{ jets} \Leftrightarrow ep \rightarrow 2 \text{ jets}$ [analytic continuation]
- $p p \rightarrow 2 \text{ jets}$: qq helicity amplitudes [JHEP 0409 (2004) 039]

✓ Ongoing effort (1997-):

- A. Gehrmann, T. Gehrmann, N. Glover, E. Remiddi, ...
- Z. Bern, L. Dixon, D. Kosower, S.Moch, P.Uwer, S. Weinzierl, ...
- Advances in mathematics: Differential eqs., nested sums
- Applications to event shapes, multijets @ 1-loop, photons

Jets (2)

Ingredients:


- $1 \rightarrow 3$ @ 2-loop: Inteference with 1 $\rightarrow$ 3 @ 0-loop ✓
- $1 \rightarrow 4$ @ 1-loop: Single soft and/or collinear regions ✓
- $1 \rightarrow 5$ @ 0-loop: Double soft ... triple collinear regions ✓

Methods:

- Optical theorem: 3-loop propagators
- Subtraction terms: Antenna functions

First (preliminary) numerical result:


- Average thrust: $\langle 1 - T \rangle = \int (1 - T) \frac{1}{\sigma_0} \frac{d\sigma}{d\Gamma} = C_F \left[\left(\frac{\alpha_s}{2\pi} \right) A + \left(\frac{\alpha_s}{2\pi} \right)^2 B + \left(\frac{\alpha_s}{2\pi} \right)^3 C + \dots \right]$
- $A = 1.57, B = 32.3, C = (-20.4 \pm 4) C_F^2 + \dots$
- A. Gehrmann, T. Gehrmann, N. Glover, LL 2004




$$\begin{aligned}
 & \frac{g^4(k_0, \epsilon_0, \mu)}{g^4(k_1, \epsilon_1, \nu)} \dots \frac{g^4(k_2, \epsilon_2, \rho)}{g^4(k_3, \epsilon_3, \sigma)} \\
 &= -i g_s^2 \lambda \left[f^{\text{tree}} f^{\text{tree}} (g^{\mu\nu} g^{\sigma\rho} - g^{\mu\sigma} g^{\nu\rho}) \right. \\
 & \quad \left. + f^{\text{tree}} f^{\text{tree}} (g^{\mu\nu} g^{\sigma\rho} - g^{\mu\sigma} g^{\nu\rho}) + f^{\text{tree}} f^{\text{tree}} (g^{\mu\sigma} g^{\nu\rho} - g^{\mu\nu} g^{\sigma\rho}) \right]
 \end{aligned}$$

Jets (3)

Thrust: Higher twist?




Diphotons: Higgs from gg?



Higgs (1)

Inclusive cross section: $gg, (bb) \rightarrow h(H, A) + X$


- R. Harlander, B. Kilgore (2002)
- C. Anastasiou, K. Melnikov (2002)
- V. Ravindran, J. Smith, W. van Neerven (2003)



Differential cross section:

- C. Anastasiou, K. Melnikov, F. Petriello (2004)

The ggh-vertex:



$$\mathcal{L} = \frac{1}{4v} C_1 Z_1 G_{\mu\nu}^a G^{a\mu\nu} H$$

$$C_1 = -\frac{1}{3\pi} \left\{ 1 + \frac{11}{4} \frac{\alpha_s}{\pi} + \left(\frac{\alpha_s}{\pi} \right)^2 \left[\frac{2777}{288} + \frac{19}{16} L_t + n_f \left(-\frac{67}{96} + \frac{1}{3} L_t \right) \right] + \mathcal{O}(\alpha_s^3) \right\}$$


$$Z_1 = 1 - \frac{\alpha_s \beta_0}{\pi \epsilon} + \left(\frac{\alpha_s}{\pi} \right)^2 \left[\frac{\beta_0^2}{\epsilon^2} - \frac{\beta_1}{\epsilon} \right] + \mathcal{O}(\alpha_s^3)$$

Ingredients:

- $gg \rightarrow h$ @ 2-loop; $gg, qq \rightarrow hg$ @ 1-loop; $gg, qq \rightarrow hgg, qq$ @ 0-loop


Higgs (2)

Forward-going b-jets:



Harlander, Kilgore, PRD 68 (2003) 013001 M_H [GeV]

Diphotons + no hard jets:



Polarized Drell-Yan Process




NNLO coefficient function:

• V. Ravindran, J. Smith, W. van Neerven (2004)

γ_5 : HVBM-scheme, evanescent counterterms

Cross section:

Asymmetry:



Heavy Quarks

$e^+e^- \rightarrow QQ$: Vertex corrections @ 2-loop


- Vector: W. Bernreuther et al., NPB 706 (2005) 245
- Axial-vector: W. Bernreuther et al., NPB 712 (2005) 229

Method:

- Two scale problem: Q, m_Q
- Laporta reduction, IBP, Lorentz invar.
- 17 master integrals: Polylogarithms
- Renormalization: $Z_m^{OS}, Z_2^{OS}; Z_g^{MS}, Z_3^{MS}$
- Confirm threshold expansion results

Still missing: Real emission

- 1-loop single-subtraction terms
- 0-loop double-subtraction terms



Twistor Methods

- ✓ New approach to multiparticle amplitudes:
 - Relation of pert. gauge theory to string instanton expansion:
 - Twistor space: Represent 4-momenta as bi-spinors
 - Motivated by $N=4$ SYM, but works for all massless theories:
 - MHV amplitudes [scalar vertices] + scalar propagators
 - F. Cachazo, P. Svrcek, E. Witten, JHEP 0409 (2004) 006
- ✓ Tree-level: Established
 - Pure gauge theory: Next-to-MHV, recursion relations
 - Processes with fermions: MHV with quarks and gluinos
- ✓ Loop-level: Cut constructibility/new CSW-diagrams
 - Works for $N = 4$ MHV @ 1-loop for up to 7-gluons
 - Extension to $N < 4$, next-to-MHV and 2-loop unclear

Summary

✓ Inclusive processes:

- NNLO calculation completed for $F_{1,2,3} \rightarrow$ first NNLO fits
- NNLO calculation completed for Higgs and DY production

✓ Less inclusive processes:

- $e^+e^- \rightarrow 3 \text{ jets}$: $1 \rightarrow 4$ subtraction terms partially completed
 - $\rightarrow$ First (preliminary and partial) result for average thrust
- $e^+e^- \rightarrow QQ$: $1 \rightarrow 2$ vector and axial-vector vertex @ 2-loop
 - $\rightarrow$ Subtraction terms still missing
- $p p \rightarrow 2 \text{ jets}$: $2 \rightarrow 2$ quark helicity amplitudes completed
 - $\rightarrow$ Subtraction terms still missing

✓ Multi-particle processes: Twistor methods

- Tree-level: Works for non-SUSY, non-MHV, also fermions
- Loop-level: Works for $N=4$ SYM @ 1-loop, extension unclear