



PHENO 2006

Mass Effects in $Wb\bar{b}$ Production at Hadron Colliders

Pheno Symposium 2006, Madison, Wisconsin

Fernando Febres Cordero

in collaboration with

Laura Reina and Doreen Wackerroth

FSU - SUNY at Buffalo

May 16, 2006




Outline

- **Motivation:** Improving theoretical predictions of QCD backgrounds
- Brief review of the **calculation** of fully differential cross sections
- **Results:** comparison with massless approximation at LO and NLO: Theoretical uncertainties and distributions
- **Summary**




Motivation

- $Wb\bar{b}$ @ hadron colliders is background to important processes:




Light SM-like Higgs boson production
signal known @ NNLO



Single top production
signal known @ NLO

- Full massive calculation is only known @ LO. NLO calculation with $mb = 0$ approximation available in a Montecarlo program (MCFM [J. Campbell and R.K Ellis]): [R.K Ellis et al hep-ph/9810489]







Precision QCD calculation

Radiative corrections of order α_s to $d\sigma$

all divergences regularized using dimensional regularization



Then, after convoluting the virtual plus real contributions with the renormalized parton distribution functions, we get the cancellation of all singularities

In the massless approximation kinematical cuts are imposed to simulate mass effects



Experimental cuts



Having a fully differential cross section allows to implement any experimental cut to calculate total cross sections and distributions. We have produced data for the **Tevatron** ($\sqrt{s} = 1.96$ TeV), with the following cuts:

- Transverse momentum of the b-quarks: $p_t > p_{t \text{ min}}$ (15 GeV) for both b and \bar{b} quarks
- Pseudorapidity: $|\eta| < \eta_{max}$ (2) for both b and \bar{b} quarks
- kt-jet algorithm with $R_{cut} = 0.7$



Check the total cross section independence of the PS-slicing parameters

soft (δ_s) and collinear (δ_c) cuts



Preliminary





Total cross section dependence in

$p_{t \text{ min}}$ and η_{max} cuts

$$(\mu = M_W + 2m_b)$$

Preliminary






Scale dependence and theoretical uncertainty

Using $\mu \in (\frac{1}{2}(M_W + 2m_b), 2(M_W + 2m_b))$

Inclusive case:
Real cross section
integrated over all PS
of the extra parton



LO \sim 30%, NLO \sim 15%


Preliminary



Scale dependence and theoretical uncertainty

Using $\mu \in (\frac{1}{2}(M_W + 2m_b), 2(M_W + 2m_b))$

Exclusive case:
Real cross section
integrated over PS
when extra parton is
not resolved





LO ~ 30%, NLO ~ 8%

Preliminary




LO and NLO distributions in the invariant mass of the $b - \bar{b}$ pair



Preliminarily, using this cuts, we have found a difference of around 7% between the massless NLO calculation and our NLO result, mostly coming from regions with small invariant mass of the $b - \bar{b}$ pair



LO and NLO comparison of massive distributions



Preliminary



Summary

- We have calculated the NLO QCD radiative correction to the production of a W with two massive jets. We observe considerable reduction of the theoretical uncertainty on the total cross section with respect to the LO calculation, allowing for better predictions of backgrounds to important processes like vector boson associated Higgs production and Single Top production.
- Mass effects are relevant only for small invariant mass of the $b - \bar{b}$ pair (below 60 GeV).
- Because of the difference with LO distribution is important to use NLO distributions for Light Higgs searches.
- We are working to produce the $Z + b\bar{b}$ counterpart of this calculation.