

A_5 Family Symmetry and the Golden Ratio Prediction for Solar Neutrino Mixing

Alexander Stuart

UW-Madison


May 12th, 2009

**Based on: L. Everett and A. Stuart, 0812.1057 [hep-ph], to appear in PRD
and work in progress**

The Standard Model

$$SU(3)_c \times SU(2)_L \times U(1)_Y$$

- Triumph of modern science, but incomplete. Predicts massless neutrinos.
- How can we mix neutrinos into our recipe?



Figuring Out the Ingredients

$$\Delta m_{ij}^2 \equiv m_i^2 - m_j^2$$


$$\Delta m_{21}^2 = 7.6_{-0.3}^{+0.5} \cdot 10^{-5} \text{ eV}^2 \text{ and}$$

$$\sin^2 \theta_{12} = 0.32_{-0.04}^{+0.05}$$

P. Huber

$$\Delta m_{31}^2 = 2.4_{-0.3}^{+0.3} \cdot 10^{-3} \text{ eV}^2 \text{ and } \sin^2 \theta_{23} = 0.5_{-0.12}^{+0.13}$$

$$\sin^2 \theta_{13} \leq 0.033$$



Generating Small Neutrino Masses

- Dirac neutrinos: add right handed neutrino to SM:

$$m_D \nu_L \overline{N}_R \quad m_{ij} = Y_{ij} \langle H \rangle$$

but very small Yukawa coupling ($\sim 10^{-12}$)

- Majorana neutrinos: seesaw mechanism
(Minkowski; Gell-Mann, Ramond, Slansky; Yanagida)

Effective Majorana neutrino mass terms (LL couplings)

$$(m_D M_{maj}^{-1} m_D^T)_{ij} L_i L_j$$

Flavor Symmetry

- Postulate a flavor symmetry (continuous or discrete) to explain mixings and masses.
- Use symmetry to forbid mass term at renormalizable level and generate mass at higher level through SSB via a flavon field.

$$T \cong A_4$$

$$\frac{1}{\sqrt{2}}$$

$$A_4 \subseteq A_5$$

$$I \cong A_5$$

$$\phi = \frac{1 + \sqrt{5}}{2}$$

$$\text{ArcTan}\left(\frac{1}{\sqrt{2}}\right) = 35.2644^\circ$$

$$\text{ArcTan}\left(\frac{1}{\phi}\right) = 31.7175^\circ$$

Y. Kajiyama, M. Raidal, and A. Strumia ($Z_2 \otimes Z_2$);
L.L. Everett and A. J. Stuart (A_5)


(Alternate golden ratio scenario: A. Adulpravitchai, A. Blum, W. Rodejohann (D_{10}))

Icosahedral Group is not crystallographic point group
so there was work to be done.

The Icosahedral Group, I

- An icosahedron is the Platonic solid that consists of 20 equilateral triangles. $\rightarrow f = 20$
- 20 triangles each have 3 sides $\rightarrow 60$ edges but 2 triangles/edge $\rightarrow 30$ edges $\rightarrow e=30$
- 20 triangles each have 3 vertices $\rightarrow 60$ vertices but 5 vertices/edge $\rightarrow v= 12$
- Are we right? $\chi(g) = 2 - 2g = v - e + f$
- I consists of all rotations that take vertices to vertices i.e. $0, \pi, \frac{2\pi}{3}, \frac{2\pi}{5}, \frac{4\pi}{5}$

$$I \cong A_5$$



May 12, 2009

Conjugacy Classes of I

Rotation by each angle forms its own conjugacy class.

Schoenflies Notation: C_n^k is a rotation by $\frac{2\pi k}{n}$

in front = # of elements in class

So for the icosahedral group we have:

$$I, 12C_5, 12C_5^2, 20C_3, 15C_2$$

Note: $1 + 12 + 12 + 15 + 20 = 60 = 1^2 + 3^2 + 3^2 + 4^2 + 5^2$ **Two triplets.**

What do we do with this information?

The Icosahedral Character Table

- A character table is a table that gives the characters of the group elements as a function of the conjugacy class.

\mathcal{I}	1	3	3'	4	5
e	1	3	3	4	5
$12C_5$	1	ϕ	$1 - \phi$	-1	0
$12C_5^2$	1	$1 - \phi$	ϕ	-1	0
$20C_3$	1	0	0	1	-1
$15C_2$	1	-1	-1	0	1

Kronecker Products of I

Use Character Table to easily obtain Kronecker Products (known)

All of this is abstract. We need actual representations.

Our work: Identified useful group presentation where golden ratio is manifest:

K. Shirai, J. Phys. Soc. Jpn. 61 2735 (1992).

Explicitly constructed C-G coefficients in Shirai basis.

$$3 \otimes 3 = 1 \oplus 3 \oplus 5$$

$$3' \otimes 3' = 1 \oplus 3' \oplus 5$$

$$3 \otimes 3' = 4 \oplus 5$$

$$3 \otimes 4 = 3' \oplus 4 \oplus 5$$

$$3' \otimes 4 = 3 \oplus 4 \oplus 5$$

$$3 \otimes 5 = 3 \oplus 3' \oplus 4 \oplus 5$$

$$3' \otimes 5 = 3 \oplus 3' \oplus 4 \oplus 5$$

$$4 \otimes 4 = 1 \oplus 3 \oplus 3' \oplus 4 \oplus 5$$

$$4 \otimes 5 = 3 \oplus 3' \oplus 4 \oplus 5 \oplus 5$$

$$5 \otimes 5 = 1 \oplus 3 \oplus 3' \oplus 4 \oplus 4 \oplus 5 \oplus 5$$

Tensor Product Decomposition

$$3 \otimes 3 \quad 3 = \begin{pmatrix} a_1 \\ a_2 \\ a_3 \end{pmatrix} \quad 3 \otimes 3'$$

$$1 = a_1 b_1 + a_2 b_2 + a_3 b_3$$

$$3 = (a_3 b_2 - b_2 a_3, a_1 b_3 - a_3 b_1, a_2 b_1 - a_1 b_2)^T$$

$$4 = \begin{pmatrix} \frac{1}{\phi} a_3 b_2 - \phi a_1 b_3 \\ \phi a_3 b_1 + \frac{1}{\phi} a_2 b_3 \\ -\frac{1}{\phi} a_1 b_1 + \phi a_2 b_2 \\ a_2 b_1 - a_1 b_2 + a_3 b_3 \end{pmatrix}$$

$$5 = \begin{pmatrix} a_2 b_2 - a_1 b_1 \\ a_2 b_1 + a_1 b_2 \\ a_3 b_2 + a_2 b_3 \\ a_1 b_3 + a_3 b_1 \\ -\frac{1}{\sqrt{3}}(a_1 b_1 + a_2 b_2 - 2a_3 b_3) \end{pmatrix}$$

$$5 = \begin{pmatrix} \frac{1}{2}(\phi^2 a_2 b_1 + \frac{1}{\phi^2} a_1 b_2 - \sqrt{5} a_3 b_3) \\ -(\phi a_1 b_1 + \frac{1}{\phi} a_2 b_2) \\ \frac{1}{\phi} a_3 b_1 - \phi a_2 b_3 \\ \phi a_3 b_2 + \frac{1}{\phi} a_1 b_3 \\ \frac{\sqrt{3}}{2}(\frac{1}{\phi} a_2 b_1 + \phi a_1 b_2 + a_3 b_3) \end{pmatrix}$$

How to Build an A_5 Flavor Model

Seek scenario with: maximal θ_{atm} , zero θ_{13} , and “golden” θ_{sol} .

Our approach: atmospheric mixing from the charged leptons, solar mixing from neutrinos

Choose to assign irreps to SM fields: $L \rightarrow 3$, $\bar{e} \rightarrow 3'$

Tree level: $L \bar{e}$ ($3 \times 3'$): vanish

LL (3×3): degenerate

$$3 \otimes 3' = 4 \oplus 5 \quad 3 \otimes 3 = 1 \oplus 3 \oplus 5$$

How to Build an A_5 Flavor Model

Flavon sector can dramatically alter tree-level pattern:
Here, take flavon fields ξ , χ , and ψ

Charge assignment: $(L, \bar{e}, \xi, \chi, \psi) \sim (3, 3', 5, 4, 5')$

The invariant effective Lagrangian is:

$$- \mathcal{L}_{mass} = \frac{\alpha_{ijk}}{MM'} L_i H L_j H \xi_k + \frac{\beta_{ijk}}{M'} L_i \bar{e}_j H \psi_k + \frac{\gamma_{ijl}}{M'} L_i \bar{e}_j H \chi_l + \text{h.c.}$$

Spontaneous Symmetry Breaking

- **Assume flavon field vevs:**

$$\begin{pmatrix} \xi_1 \\ \xi_2 \\ \xi_3 \\ \xi_4 \\ \xi_5 \end{pmatrix} \longrightarrow \frac{\sqrt{3}}{2\alpha} \begin{pmatrix} \frac{1}{\sqrt{15}}(m_2 - m_1) \\ \frac{2}{\sqrt{15}}(m_2 - m_1) \\ 0 \\ 0 \\ -(m_1 + m_2) \end{pmatrix} \quad \begin{pmatrix} \psi_1 \\ \psi_2 \\ \psi_3 \\ \psi_4 \\ \psi_5 \end{pmatrix} \longrightarrow \frac{1}{2\sqrt{6}\beta} \begin{pmatrix} -\sqrt{\frac{5}{3}}m_\tau \\ \frac{2}{\sqrt{3}}(-\phi\sqrt{2}m_e - \frac{1}{\phi}m_\mu) \\ -\frac{2}{\sqrt{3}}\phi m_\tau \\ -\frac{2}{\sqrt{3}}\phi m_\mu \\ m_\tau \end{pmatrix}$$

$$\begin{pmatrix} \chi_1 \\ \chi_2 \\ \chi_3 \\ \chi_4 \end{pmatrix} \longrightarrow \frac{1}{3\sqrt{2}\gamma} \begin{pmatrix} -\frac{1}{\phi}m_\mu \\ \frac{1}{\phi}m_\tau \\ -\frac{\sqrt{2}}{\phi}m_e + \phi m_\mu \\ m_\tau \end{pmatrix}$$

Mass Matrices

- After spontaneous symmetry breaking, the mass matrices are:

$$\text{Neutrinos: } M_\nu = \frac{1}{\sqrt{5}} \begin{pmatrix} \phi m_1 + \frac{1}{\phi} m_2 & m_2 - m_1 & 0 \\ m_2 - m_1 & \frac{1}{\phi} m_1 + \phi m_2 & 0 \\ 0 & 0 & -\sqrt{5}(m_1 + m_2) \end{pmatrix}$$

$$\text{Charged leptons: } M_e = \frac{1}{\sqrt{2}} \begin{pmatrix} \sqrt{2} m_e & 0 & 0 \\ 0 & m_\mu & m_\tau \\ 0 & -m_\mu & m_\tau \end{pmatrix}$$

The MNSP Matrix

- Recall: $U_{MNSP} = U_{-1}U_{\nu}^{\dagger}$ where U_{-1} and U_{ν} are unitary matrices that diagonalize the left-handed fields. By construction, we have

$$U = U_{MNSP} = U_e U_{\nu}^{\dagger} = \begin{pmatrix} \sqrt{\frac{\phi}{\sqrt{5}}} & \sqrt{\frac{1}{\sqrt{5}\phi}} & 0 \\ -\frac{1}{\sqrt{2}}\sqrt{\frac{1}{\sqrt{5}\phi}} & \frac{1}{\sqrt{2}}\sqrt{\frac{\phi}{\sqrt{5}}} & -\frac{1}{\sqrt{2}} \\ -\frac{1}{\sqrt{2}}\sqrt{\frac{1}{\sqrt{5}\phi}} & \frac{1}{\sqrt{2}}\sqrt{\frac{\phi}{\sqrt{5}}} & \frac{1}{\sqrt{2}} \end{pmatrix} \mathcal{P}$$

$$\mathcal{P} = \text{Diag}(1, 1, i)$$

Features of the Model

- By design, we have $\theta_{13} = 0^\circ$ $\theta_{atm} = 45^\circ$
 $\theta_{sol} = \text{ArcTan}\left(\frac{1}{\phi}\right) = 31.7175^\circ$

Predicts a normal hierarchy with: $m_3 = m_1 + m_2$

$0\nu\beta\beta$ decay: $m_{\beta\beta} = \frac{m_1\phi}{\sqrt{5}} + \frac{m_2}{\phi\sqrt{5}}$

- These results are well within the measurements.
The solar angle is 2σ below best fit.

Outlook and Conclusion

- Summary:
 - Icosahedral (A_5) symmetry provides a rich setting for investigating the flavor puzzle.
 - “Golden Ratio” models: intriguing alternative to tri-bimaximal mixing scenarios. Virtually unexplored!
- Where to next?
 - Analyze the flavon sector dynamics (in progress).
 - Study other lepton mixing scenarios, investigate quark sector, SUSY embeddings,...