

Interpretation of Higgs data within the SMEFT

Ken Mimasu

King's College London

14th Conference on the Intersections of Particle and Nuclear Physics

30th August 2022



[J. Ellis, M. Madigan, KM, V. Sanz & T. You; JHEP 04 (2021) 279]

fitmaker <https://gitlab.com/kenmimasu/fitrepo>

Where are we?

10 years since the Higgs discovery

- No clear sign of new physics at the TeV scale
- Direct searches are saturating the energy frontier



[CERN-EP-2020-049]

ATLAS Exotics Searches* - 95% CL Upper Exclusion Limits

Status: May 2020

$\int \mathcal{L} dt = (3.2 - 139) \text{ fb}^{-1}$

ATLAS Preliminary

$\sqrt{s} = 8, 13 \text{ TeV}$

Model	ℓ, γ	Jets [†]	E_T^{miss}	$\int \mathcal{L} dt [\text{fb}^{-1}]$	Limit	Reference
Extra dimensions						
ADD $G_{KK} + g/q$	$0 e, \mu$	1-4 j	Yes	36.1	M_D 7.7 TeV	$n=2$
ADD non-resonant $\gamma\gamma$	2γ	-	-	36.7	M_S 8.6 TeV	$n=3$ HLZ NLO
ADD QBH	-	2j	-	37.0	M_{th} 8.9 TeV	$n=6$
ADD BH high Σp_T	$\geq 1 e, \mu$	$\geq 2 j$	-	3.2	M_{th} 8.2 TeV	$n=6, M_D = 3 \text{ TeV, rot BH}$
ADD BH multijet	-	$\geq 3 j$	-	3.6	M_{th} 9.55 TeV	$n=6, M_D = 3 \text{ TeV, rot BH}$
RS1 $G_{KK} \rightarrow \gamma\gamma$	2γ	-	-	36.7	$G_{KK} \text{ mass}$ 4.1 TeV	$k/\overline{M}_{\text{Pl}} = 0.1$
Bulk RS $G_{KK} \rightarrow WW/ZZ$	multi-channel	-	-	36.1	$G_{KK} \text{ mass}$ 2.3 TeV	$k/\overline{M}_{\text{Pl}} = 1.0$
Bulk RS $G_{KK} \rightarrow WV \rightarrow \ell\nu q\bar{q}$	$1 e, \mu$	2j/1J	Yes	139	$G_{KK} \text{ mass}$ 2.0 TeV	$k/\overline{M}_{\text{Pl}} = 1.0$
Bulk RS $G_{KK} \rightarrow t\bar{t}$	$1 e, \mu$	$\geq 1 b, \geq 1J/2j$	Yes	36.1	$G_{KK} \text{ mass}$ 3.8 TeV	$\Gamma/m = 15\%$
2UED / RPP	$1 e, \mu$	$\geq 2 b, \geq 3 j$	Yes	36.1	$KK \text{ mass}$ 1.8 TeV	Tier (1,1), $\mathcal{B}(A^{(1,1)} \rightarrow t\bar{t}) = 1$
Gauge bosons						
SSM $Z' \rightarrow \ell\ell$	$2 e, \mu$	-	-	139	$Z' \text{ mass}$ 5.1 TeV	
SSM $Z' \rightarrow \tau\tau$	2τ	-	-	36.1	$Z' \text{ mass}$ 2.42 TeV	
Leptophobic $Z' \rightarrow b\bar{b}$	-	2b	-	36.1	$Z' \text{ mass}$ 2.1 TeV	
Leptophobic $Z' \rightarrow t\bar{t}$	$0 e, \mu$	$\geq 1 b, \geq 2 J$	Yes	139	$Z' \text{ mass}$ 4.1 TeV	$\Gamma/m = 1.2\%$
SSM $W' \rightarrow \ell\nu$	$1 e, \mu$	-	Yes	139	$W' \text{ mass}$ 6.0 TeV	
SSM $W' \rightarrow \tau\nu$	1τ	-	Yes	36.1	$W' \text{ mass}$ 3.7 TeV	
HVT $W' \rightarrow WZ \rightarrow \ell\nu q\bar{q}$ model B	$1 e, \mu$	2j/1J	Yes	139	$W' \text{ mass}$ 4.0 TeV	$g_V = 3$
HVT $V' \rightarrow WV \rightarrow qq\bar{q}\bar{q}$ model B	$0 e, \mu$	2J	-	139	$V' \text{ mass}$ 3.8 TeV	$g_V = 3$
HVT $V' \rightarrow WH/ZH$ model B	multi-channel	-	-	36.1	$V' \text{ mass}$ 2.93 TeV	$g_V = 3$
HVT $W' \rightarrow WH$ model B	$0 e, \mu$	$\geq 1 b, \geq 2 J$	-	139	$W' \text{ mass}$ 3.2 TeV	$g_V = 3$
LRSM $W_R \rightarrow t\bar{b}$	multi-channel	-	-	36.1	$W_R \text{ mass}$ 3.25 TeV	
LRSM $W_R \rightarrow \mu N_R$	2μ	1J	-	80	$W_R \text{ mass}$ 5.0 TeV	$m(N_R) = 0.5 \text{ TeV, } g_L = g_R$
CI						
CI $qq\bar{q}\bar{q}$	-	2j	-	37.0	Λ 21.8 TeV	η_{LL}
CI $\ell\ell q\bar{q}$	$2 e, \mu$	-	-	139	Λ 35.8 TeV	η_{LL}
CI $t\bar{t}t\bar{t}$	$\geq 1 e, \mu$	$\geq 1 b, \geq 1 j$	Yes	36.1	Λ 2.57 TeV	$ C_{41} = 4\pi$
DM						
Axial-vector mediator (Dirac DM)	$0 e, \mu$	1-4 j	Yes	36.1	m_{med} 1.65 TeV	$g_q = 0.25, g_\ell = 1.0, m(\chi) = 1 \text{ GeV}$
Colored scalar mediator (Dirac DM)	$0 e, \mu$	1-4 j	Yes	36.1	m_{med} 1.67 TeV	$g = 1.0, m(\chi) = 1 \text{ GeV}$
$VV\chi\chi$ EFT (Dirac DM)	$0 e, \mu$	1j, $\leq 1 j$	Yes	3.2	M_χ 700 GeV	$m(\chi) < 150 \text{ GeV}$
Scalar reson. $\phi \rightarrow t\bar{t}$ (Dirac DM)	$0-1 e, \mu$	1b, 0-1J	Yes	36.1	m_ϕ 3.4 TeV	$y = 0.4, \lambda = 0.2, m(\chi) = 10 \text{ GeV}$
LQ						
Scalar LQ 1 st gen	$1, 2 e$	$\geq 2 j$	Yes	36.1	$LQ \text{ mass}$ 1.4 TeV	$\beta = 1$
Scalar LQ 2 nd gen	$1, 2 e$	$\geq 2 j$	Yes	36.1	$LQ \text{ mass}$ 1.56 TeV	$\beta = 1$
Scalar LQ 3 rd gen	2τ	2b	-	36.1	$LQ \text{ mass}$ 1.03 TeV	$\mathcal{B}(LQ_3^+ \rightarrow b\tau) = 1$
Scalar LQ 3 rd gen	$0-1 e, \mu$	2b	Yes	36.1	$LQ \text{ mass}$ 970 GeV	$\mathcal{B}(LQ_3^+ \rightarrow t\tau) = 0$
Heavy quarks						
VLQ $TT \rightarrow Ht/Zt/Wb + X$	multi-channel	-	-	36.1	$T \text{ mass}$ 1.3 TeV	SU(2) doublet
VLQ $BB \rightarrow Wt/Zb + X$	multi-channel	-	-	36.1	$B \text{ mass}$ 1.34 TeV	SU(2) doublet
VLQ $T_{5/3} T_{5/3} T_{5/3} \rightarrow Wt + X$	2(SS) $\geq 3 e, \mu \geq 1 b, \geq 1 j$	Yes	36.1	$T_{5/3} \text{ mass}$ 1.64 TeV	$\mathcal{B}(T_{5/3} \rightarrow Wt) = 1, c(T_{5/3} Wt) = 1$	
VLQ $Y \rightarrow Wb + X$	$1 e, \mu$	$\geq 1 b, \geq 1 j$	Yes	36.1	$Y \text{ mass}$ 1.85 TeV	$\mathcal{B}(Y \rightarrow Wb) = 1, c_R(Wb) = 1$
VLQ $B \rightarrow Hb + X$	$0 e, \mu, 2 \gamma$	$\geq 1 b, \geq 1 j$	Yes	79.8	$B \text{ mass}$ 1.21 TeV	$x_B = 0.5$
VLQ $QQ \rightarrow WqWq$	$1 e, \mu$	$\geq 4 j$	Yes	20.3	$Q \text{ mass}$ 690 GeV	
Excited fermions						
Excited quark $q^* \rightarrow qg$	-	2j	-	139	$q^* \text{ mass}$ 6.7 TeV	only u^* and d^* , $\Lambda = m(q^*)$
Excited quark $q^* \rightarrow q\gamma$	1γ	1j	-	36.7	$q^* \text{ mass}$ 5.3 TeV	only u^* and d^* , $\Lambda = m(q^*)$
Excited quark $b^* \rightarrow b\gamma$	-	1b, 1j	-	36.1	$b^* \text{ mass}$ 2.6 TeV	
Excited lepton ℓ^*	$3 e, \mu$	-	-	20.3	$\ell^* \text{ mass}$ 3.0 TeV	$\Lambda = 3.0 \text{ TeV}$
Excited lepton ν^*	$3 e, \mu, \tau$	-	-	20.3	$\nu^* \text{ mass}$ 1.6 TeV	$\Lambda = 1.6 \text{ TeV}$
Other						
Type III Seesaw	$1 e, \mu$	$\geq 2 j$	Yes	79.8	$N^0 \text{ mass}$ 560 GeV	
LRSM Majorana ν	2μ	2j	-	36.1	$N_R \text{ mass}$ 3.2 TeV	$m(W_R) = 4.1 \text{ TeV, } g_L = g_R$
Higgs triplet $H^{\pm\pm} \rightarrow \ell\ell$	$2, 3, 4 e, \mu$ (SS)	-	-	36.1	$H^{\pm\pm} \text{ mass}$ 870 GeV	DY production
Higgs triplet $H^{\pm\pm} \rightarrow \ell\tau$	$3 e, \mu, \tau$	-	-	20.3	$H^{\pm\pm} \text{ mass}$ 400 GeV	DY production, $\mathcal{B}(H^{\pm\pm} \rightarrow \ell\tau) = 1$
Multi-charged particles	-	-	-	36.1	multi-charged particle mass 1.22 TeV	DY production, $ q = 5e$
Magnetic monopoles	-	-	-	34.4	monopole mass 2.37 TeV	DY production, $ g = 1g_D, \text{spin } 1/2$

$\sqrt{s} = 8 \text{ TeV}$ $\sqrt{s} = 13 \text{ TeV}$ partial data $\sqrt{s} = 13 \text{ TeV}$ full data

[TeV] : 1 3 5 10


Energy & precision

Paradigm shift at the energy
frontier for BSM searches

Energy & precision

Paradigm shift at the energy frontier for BSM searches

Direct (bumps)




Energy & precision

Paradigm shift at the energy frontier for BSM searches

Direct (bumps)

Indirect (tails)




Energy & precision

Paradigm shift at the energy frontier for BSM searches

Direct (bumps)

Indirect (tails)

⇒ New physics is heavy



Energy & precision

Paradigm shift at the energy frontier for BSM searches

Direct (bumps)


Indirect (tails)

⇒ New physics is heavy

Heavy new physics

Precision measurements

High energy



Energy & precision

Paradigm shift at the energy frontier for BSM searches

Direct (bumps)



Indirect (tails)

⇒ New physics is heavy

Heavy new physics

Precision measurements

High energy



**Standard Model
Effective Field Theory
(SMEFT)**


Energy & precision

Paradigm shift at the energy frontier for BSM searches

Direct (bumps)

Indirect (tails)


⇒ New physics is heavy



Heavy new physics

Precision measurements

High energy



**Standard Model
Effective Field Theory
(SMEFT)**

A QFT parameter space for BSM interactions between SM particles

SMEFT: SM v2.0

$$\mathcal{L}_{\text{eff}} = \sum_i \frac{c_i \mathcal{O}_i^D}{\Lambda^{D-4}}$$

SM is low energy effective description

- Supplemented by a tower of irrelevant operators
- Respecting low energy field content & symmetries

$$\text{SU}(3)_c \times \text{SU}(2)_L \times \text{U}(1)_Y$$

$$\varphi = \begin{pmatrix} G^+ \\ v + h + iG^0 \end{pmatrix} : \mathbf{2}_{\frac{1}{2}}$$

SMEFT: SM v2.0

$$\mathcal{L}_{\text{eff}} = \sum_i \frac{c_i \mathcal{O}_i^D}{\Lambda^{D-4}}$$

SM is low energy effective description

- Supplemented by a tower of irrelevant operators
- Respecting low energy field content & symmetries

$$\text{SU}(3)_c \times \text{SU}(2)_L \times \text{U}(1)_Y$$

$$\varphi = \begin{pmatrix} G^+ \\ v + h + iG^0 \end{pmatrix} : \mathbf{2}_{\frac{1}{2}}$$

aTGC

$$X^3 : \epsilon_{IJK} W_{\mu\nu}^I W^{J,\nu\rho} W_{\rho}^{K,\mu}$$

$$X^2 H^2 : (\varphi^\dagger \varphi)^2 G_{\mu\nu}^a G_a^{\mu\nu}$$

ggh(h)

λ_h

$$H^6 : (\varphi^\dagger \varphi)^3$$

$$H^4 D^2 : (\varphi^\dagger D^\mu \varphi)^* (\varphi^\dagger D^\mu \varphi)$$

δM_Z

y_f

$$\psi^2 H^3 : (\varphi^\dagger \varphi)^2 (\bar{q}_i u_j \tilde{\varphi})$$

$$\psi^2 XH : (\bar{q}_i \sigma^{\mu\nu} u_j \tilde{\varphi}) B_{\mu\nu}$$

'dipole'

ffV

$$\psi^2 H^2 D : (\varphi^\dagger \overleftrightarrow{D}_\mu \varphi) (\bar{q}_i \gamma^\mu q_j)$$

$$\psi^4 : (\bar{q}_i \gamma^\mu q_j) (\bar{q}_k \gamma_\mu q_l)$$

4F

SMEFT: SM v2.0

$$\mathcal{L}_{\text{eff}} = \sum_i \frac{c_i \mathcal{O}_i^D}{\Lambda^{D-4}}$$

SM is low energy effective description

- Supplemented by a tower of irrelevant operators
- Respecting low energy field content & symmetries

$$\text{SU}(3)_c \times \text{SU}(2)_L \times \text{U}(1)_Y$$

$$\varphi = \begin{pmatrix} G^+ \\ v + h + iG^0 \end{pmatrix} : \mathbf{2}_{\frac{1}{2}}$$

aTGC

$$X^3 : \epsilon_{IJK} W_{\mu\nu}^I W^{J,\nu\rho} W_{\rho}^{K,\mu}$$

$$X^2 H^2 : (\varphi^\dagger \varphi)^2 G_{\mu\nu}^a G_a^{\mu\nu}$$

ggh(h)

λ_h

$$H^6 : (\varphi^\dagger \varphi)^3$$

$$H^4 D^2 : (\varphi^\dagger D^\mu \varphi)^* (\varphi^\dagger D^\mu \varphi)$$

δM_Z

y_f

$$\psi^2 H^3 : (\varphi^\dagger \varphi)^2 (\bar{q}_i u_j \tilde{\varphi})$$

$$\psi^2 XH : (\bar{q}_i \sigma^{\mu\nu} u_j \tilde{\varphi}) B_{\mu\nu}$$

'dipole'

ffV

$$\psi^2 H^2 D : (\varphi^\dagger \overleftrightarrow{D}_\mu \varphi) (\bar{q}_i \gamma^\mu q_j)$$

$$\psi^4 : (\bar{q}_i \gamma^\mu q_j) (\bar{q}_k \gamma_\mu q_l)$$

4F

More than 'just' a parametrisation of ignorance

- **Unlike** anomalous couplings
- **Renormalisable** QFT (order-by-order)
- **Finite energy range** ($\sim \Lambda$)
- Well defined **matching** procedure

SMEFT is...

$$\mathcal{L}_{\text{eff}} = \sum_i \frac{c_i \mathcal{O}_i^D}{\Lambda^{D-4}}$$

SMEFT is...

$$\mathcal{L}_{\text{eff}} = \sum_i \frac{c_i \mathcal{O}_i^D}{\Lambda^{D-4}}$$

Model independent

- Underlying assumptions

*Heavy new physics: $M > E_{\text{exp}}$
SM field content & gauge symmetries
Linear EWSB: Higgs = doublet*

SMEFT is...

$$\mathcal{L}_{\text{eff}} = \sum_i \frac{c_i \mathcal{O}_i^D}{\Lambda^{D-4}}$$

Model independent

- Underlying assumptions

Heavy new physics: $M > E_{\text{exp}}$

SM field content & gauge symmetries

Linear EWSB: Higgs = doublet

Systematically improvable

- Double expansion

higher dim. $\frac{E^2}{\Lambda^2}$ & $\{g_s, g, g'\}$ *more loops*

SMEFT is...

$$\mathcal{L}_{\text{eff}} = \sum_i \frac{c_i \mathcal{O}_i^D}{\Lambda^{D-4}}$$

Model independent

- Underlying assumptions

*Heavy new physics: $M > E_{\text{exp}}$
SM field content & gauge symmetries
Linear EWSB: Higgs = doublet*

Systematically improvable

- Double expansion *higher dim.* $\frac{E^2}{\Lambda^2}$ & $\{g_s, g, g'\}$ *more loops*

Global

- *Model independence*: we don't know what operators NP will generate
- *Patterns & correlations* among operators & observables are key

SMEFT is...

$$\mathcal{L}_{\text{eff}} = \sum_i \frac{c_i \mathcal{O}_i^D}{\Lambda^{D-4}}$$

Model independent

- Underlying assumptions

*Heavy new physics: $M > E_{\text{exp}}$
SM field content & gauge symmetries
Linear EWSB: Higgs = doublet*

Systematically improvable

- Double expansion *higher dim.* $\frac{E^2}{\Lambda^2}$ & $\{g_s, g, g'\}$ *more loops*

Global

- *Model independence*: we don't know what operators NP will generate
- *Patterns & correlations* among operators & observables are key
- *Ultimate goal*: complete SMEFT likelihood confronted with HEP data

EWPO, Higgs, multiboson, top, DY, flavor, ...

SMEFT is...

$$\mathcal{L}_{\text{eff}} = \sum_i \frac{c_i \mathcal{O}_i^D}{\Lambda^{D-4}}$$

Model independent

- Underlying assumptions

*Heavy new physics: $M > E_{\text{exp}}$
SM field content & gauge symmetries
Linear EWSB: Higgs = doublet*

Systematically improvable

- Double expansion *higher dim.* $\frac{E^2}{\Lambda^2}$ & $\{g_s, g, g'\}$ *more loops*

Global

- *Model independence*: we don't know what operators NP will generate
- *Patterns & correlations* among operators & observables are key
- *Ultimate goal*: complete SMEFT likelihood confronted with HEP data

EWPO, Higgs, multiboson, top, DY, flavor, ...

Established part of LHC programme

SMEFT interpretation

Ingredients:

$$\Delta o_n = o_n^{\text{EXP}} - o_n^{\text{SM}} = \sum_i \frac{a_{n,i}^{(6)}(\mu) c_i^{(6)}(\mu)}{\Lambda^2} + \mathcal{O}\left(\frac{1}{\Lambda^3}\right)$$

SMEFT interpretation

Ingredients:

$$\boxed{\Delta o_n} = o_n^{\text{EXP}} - o_n^{\text{SM}} = \sum_i \frac{a_{n,i}^{(6)}(\mu) c_i^{(6)}(\mu)}{\Lambda^2} + \mathcal{O}\left(\frac{1}{\Lambda^3}\right)$$

Global nature

As many observables
as possible

Identify patterns &
correlations in fits

Exploit energy-growth

SMEFT interpretation

Ingredients:

$$\Delta o_n = o_n^{\text{EXP}} - o_n^{\text{SM}} = \sum_i \frac{a_{n,i}^{(6)}(\mu) c_i^{(6)}(\mu)}{\Lambda^2} + \mathcal{O}\left(\frac{1}{\Lambda^3}\right)$$

Global nature

As many observables as possible

Identify patterns & correlations in fits

Exploit energy-growth

Sensitivity

Experiment:

Best measurements & understanding of uncertainties and correlations

Theory:

Best available predictions for observables (NLO, NNLO, N3LO,...)

SMEFT interpretation

Ingredients:

$$\Delta o_n = o_n^{\text{EXP}} - o_n^{\text{SM}} = \sum_i \frac{a_{n,i}^{(6)}(\mu) c_i^{(6)}(\mu)}{\Lambda^2} + \mathcal{O}\left(\frac{1}{\Lambda^3}\right)$$

Global nature

As many observables as possible

Identify patterns & correlations in fits

Exploit energy-growth

Sensitivity

Experiment:

Best measurements & understanding of uncertainties and correlations

Theory:

Best available predictions for observables (NLO, NNLO, N3LO,...)

Interpretation

Relies on accurate knowledge of the size & correlation among a_i



Determining $c_i^{(6)}$ requires most precise available SMEFT predictions

Higgs data

Signal strengths, $\mu_{i \rightarrow f} \sim \frac{\sigma_{i \rightarrow H} \times \text{BR}_{H \rightarrow f}}{\sigma_{i \rightarrow H}^{\text{SM}} \times \text{BR}_{H \rightarrow f}^{\text{SM}}}$

κ -framework interpretation

CMS [CMS; Nature 607 (2022) 7917, 60-68] 138 fb⁻¹ (13 TeV)



+ correlations


$\mu_{\text{tot}} = 1.002 \pm 0.057$

$\mu_{i \rightarrow f} \propto \kappa_i^2 \kappa_f^2$

Higgs data

Simplified Template Cross Sections (STXS)

[ATLAS; Nature 607 (2022) 7917, 52-59]



- Production binned in Higgs/associated particle kinematics (unfolded to SM)

$ggF : \{p_T^H, n_{\text{jet}}, m_{jj}, \dots\}$ $EW qqH : \{p_T^H, n_{\text{jet}}, m_{jj}, \dots\}$ $VH \rightarrow \text{leptons} : \{p_T^V, n_{\text{jet}}, \dots\}$ $t\bar{t}H : \{p_T^H, \dots\}$


- Measured for each decay mode: $b\bar{b}, \gamma\gamma, \tau\tau, 4\ell, \dots$

- Detailed correlation information for consistent combination


Higgs data

Differential cross sections

[CMS; PRL 128 (2022) 081805]



[ATLAS; JHEP 08 (2022) 027]




- Most fine-grained binning → best individual sensitivity
- No correlation information with other channels
- Statistical overlap with, e.g., STXS

Higgs data


CP properties

[ATLAS; PRL 125 (2020) 061802]



Rare decays

[CMS-HIG-19-014]



Breadth of SMEFT framework

- All measurements can be interpreted in a **global** analysis
- Data \Leftrightarrow Wilson coefficients: search of hints of heavy new physics

Bosonic operators

[Grzadkowski et al.; JHEP 10 (2010) 085]


X^3		φ^6 and $\varphi^4 D^2$		$\psi^2 \varphi^3$	
Q_G	$f^{ABC} G_\mu^{A\nu} G_\nu^{B\rho} G_\rho^{C\mu}$	Q_φ	$(\varphi^\dagger \varphi)^3$	$Q_{e\varphi}$	$(\varphi^\dagger \varphi)(\bar{l}_p e_r \varphi)$
$Q_{\tilde{G}}$	$f^{ABC} \tilde{G}_\mu^{A\nu} G_\nu^{B\rho} G_\rho^{C\mu}$	$Q_{\varphi\Box}$	$(\varphi^\dagger \varphi)\Box(\varphi^\dagger \varphi)$	$Q_{u\varphi}$	$(\varphi^\dagger \varphi)(\bar{q}_p u_r \tilde{\varphi})$
Q_W	$\varepsilon^{IJK} W_\mu^{I\nu} W_\nu^{J\rho} W_\rho^{K\mu}$	$Q_{\varphi D}$	$(\varphi^\dagger D^\mu \varphi)^* (\varphi^\dagger D_\mu \varphi)$	$Q_{d\varphi}$	$(\varphi^\dagger \varphi)(\bar{q}_p d_r \varphi)$
$Q_{\tilde{W}}$	$\varepsilon^{IJK} \tilde{W}_\mu^{I\nu} W_\nu^{J\rho} W_\rho^{K\mu}$				
$X^2 \varphi^2$		$\psi^2 X \varphi$		$\psi^2 \varphi^2 D$	
$Q_{\varphi G}$	$\varphi^\dagger \varphi G_{\mu\nu}^A G^{A\mu\nu}$	Q_{eW}	$(\bar{l}_p \sigma^{\mu\nu} e_r) \tau^I \varphi W_{\mu\nu}^I$	$Q_{\varphi l}^{(1)}$	$(\varphi^\dagger i \overleftrightarrow{D}_\mu \varphi)(\bar{l}_p \gamma^\mu l_r)$
$Q_{\varphi \tilde{G}}$	$\varphi^\dagger \varphi \tilde{G}_{\mu\nu}^A G^{A\mu\nu}$	Q_{eB}	$(\bar{l}_p \sigma^{\mu\nu} e_r) \varphi B_{\mu\nu}$	$Q_{\varphi l}^{(3)}$	$(\varphi^\dagger i \overleftrightarrow{D}_\mu^I \varphi)(\bar{l}_p \tau^I \gamma^\mu l_r)$
$Q_{\varphi W}$	$\varphi^\dagger \varphi W_{\mu\nu}^I W^{I\mu\nu}$	Q_{uG}	$(\bar{q}_p \sigma^{\mu\nu} T^A u_r) \tilde{\varphi} G_{\mu\nu}^A$	$Q_{\varphi e}$	$(\varphi^\dagger i \overleftrightarrow{D}_\mu \varphi)(\bar{e}_p \gamma^\mu e_r)$
$Q_{\varphi \tilde{W}}$	$\varphi^\dagger \varphi \tilde{W}_{\mu\nu}^I W^{I\mu\nu}$	Q_{uW}	$(\bar{q}_p \sigma^{\mu\nu} u_r) \tau^I \tilde{\varphi} W_{\mu\nu}^I$	$Q_{\varphi q}^{(1)}$	$(\varphi^\dagger i \overleftrightarrow{D}_\mu \varphi)(\bar{q}_p \gamma^\mu q_r)$
$Q_{\varphi B}$	$\varphi^\dagger \varphi B_{\mu\nu} B^{\mu\nu}$	Q_{uB}	$(\bar{q}_p \sigma^{\mu\nu} u_r) \tilde{\varphi} B_{\mu\nu}$	$Q_{\varphi q}^{(3)}$	$(\varphi^\dagger i \overleftrightarrow{D}_\mu^I \varphi)(\bar{q}_p \tau^I \gamma^\mu q_r)$
$Q_{\varphi \tilde{B}}$	$\varphi^\dagger \varphi \tilde{B}_{\mu\nu} B^{\mu\nu}$	Q_{dG}	$(\bar{q}_p \sigma^{\mu\nu} T^A d_r) \varphi G_{\mu\nu}^A$	$Q_{\varphi u}$	$(\varphi^\dagger i \overleftrightarrow{D}_\mu \varphi)(\bar{u}_p \gamma^\mu u_r)$
$Q_{\varphi WB}$	$\varphi^\dagger \tau^I \varphi W_{\mu\nu}^I B^{\mu\nu}$	Q_{dW}	$(\bar{q}_p \sigma^{\mu\nu} d_r) \tau^I \varphi W_{\mu\nu}^I$	$Q_{\varphi d}$	$(\varphi^\dagger i \overleftrightarrow{D}_\mu \varphi)(\bar{d}_p \gamma^\mu d_r)$
$Q_{\varphi \tilde{W}B}$	$\varphi^\dagger \tau^I \varphi \tilde{W}_{\mu\nu}^I B^{\mu\nu}$	Q_{dB}	$(\bar{q}_p \sigma^{\mu\nu} d_r) \varphi B_{\mu\nu}$	$Q_{\varphi ud}$	$i(\tilde{\varphi}^\dagger D_\mu \varphi)(\bar{u}_p \gamma^\mu d_r)$

Bosonic operators

$\phi^n D^m$

[Grzadkowski et al.; JHEP 10 (2010) 085]

X^3		φ^6 and $\varphi^4 D^2$		$\psi^2 \varphi^3$	
Q_G	$f^{ABC} G_\mu^{A\nu} G_\nu^{B\rho} G_\rho^{C\mu}$	Q_φ	$(\varphi^\dagger \varphi)^3$	$Q_{e\varphi}$	$(\varphi^\dagger \varphi)(\bar{l}_p e_r \varphi)$
$Q_{\tilde{G}}$	$f^{ABC} \tilde{G}_\mu^{A\nu} G_\nu^{B\rho} G_\rho^{C\mu}$	$Q_{\varphi\Box}$	$(\varphi^\dagger \varphi)\Box(\varphi^\dagger \varphi)$	$Q_{u\varphi}$	$(\varphi^\dagger \varphi)(\bar{q}_p u_r \tilde{\varphi})$
Q_W	$\varepsilon^{IJK} W_\mu^{I\nu} W_\nu^{J\rho} W_\rho^{K\mu}$	$Q_{\varphi D}$	$(\varphi^\dagger D^\mu \varphi)^* (\varphi^\dagger D_\mu \varphi)$	$Q_{d\varphi}$	$(\varphi^\dagger \varphi)(\bar{q}_p d_r \varphi)$
$Q_{\tilde{W}}$	$\varepsilon^{IJK} \tilde{W}_\mu^{I\nu} W_\nu^{J\rho} W_\rho^{K\mu}$				
$X^2 \varphi^2$		$\psi^2 X \varphi$		$\psi^2 \varphi^2 D$	
$Q_{\varphi G}$	$\varphi^\dagger \varphi G_{\mu\nu}^A G^{A\mu\nu}$	Q_{eW}	$(\bar{l}_p \sigma^{\mu\nu} e_r) \tau^I \varphi W_{\mu\nu}^I$	$Q_{\varphi l}^{(1)}$	$(\varphi^\dagger i \overleftrightarrow{D}_\mu \varphi)(\bar{l}_p \gamma^\mu l_r)$
$Q_{\varphi \tilde{G}}$	$\varphi^\dagger \varphi \tilde{G}_{\mu\nu}^A G^{A\mu\nu}$	Q_{eB}	$(\bar{l}_p \sigma^{\mu\nu} e_r) \varphi B_{\mu\nu}$	$Q_{\varphi l}^{(3)}$	$(\varphi^\dagger i \overleftrightarrow{D}_\mu^I \varphi)(\bar{l}_p \tau^I \gamma^\mu l_r)$
$Q_{\varphi W}$	$\varphi^\dagger \varphi W_{\mu\nu}^I W^{I\mu\nu}$	Q_{uG}	$(\bar{q}_p \sigma^{\mu\nu} T^A u_r) \tilde{\varphi} G_{\mu\nu}^A$	$Q_{\varphi e}$	$(\varphi^\dagger i \overleftrightarrow{D}_\mu \varphi)(\bar{e}_p \gamma^\mu e_r)$
$Q_{\varphi \tilde{W}}$	$\varphi^\dagger \varphi \tilde{W}_{\mu\nu}^I W^{I\mu\nu}$	Q_{uW}	$(\bar{q}_p \sigma^{\mu\nu} u_r) \tau^I \tilde{\varphi} W_{\mu\nu}^I$	$Q_{\varphi q}^{(1)}$	$(\varphi^\dagger i \overleftrightarrow{D}_\mu \varphi)(\bar{q}_p \gamma^\mu q_r)$
$Q_{\varphi B}$	$\varphi^\dagger \varphi B_{\mu\nu} B^{\mu\nu}$	Q_{uB}	$(\bar{q}_p \sigma^{\mu\nu} u_r) \tilde{\varphi} B_{\mu\nu}$	$Q_{\varphi q}^{(3)}$	$(\varphi^\dagger i \overleftrightarrow{D}_\mu^I \varphi)(\bar{q}_p \tau^I \gamma^\mu q_r)$
$Q_{\varphi \tilde{B}}$	$\varphi^\dagger \varphi \tilde{B}_{\mu\nu} B^{\mu\nu}$	Q_{dG}	$(\bar{q}_p \sigma^{\mu\nu} T^A d_r) \varphi G_{\mu\nu}^A$	$Q_{\varphi u}$	$(\varphi^\dagger i \overleftrightarrow{D}_\mu \varphi)(\bar{u}_p \gamma^\mu u_r)$
$Q_{\varphi WB}$	$\varphi^\dagger \tau^I \varphi W_{\mu\nu}^I B^{\mu\nu}$	Q_{dW}	$(\bar{q}_p \sigma^{\mu\nu} d_r) \tau^I \varphi W_{\mu\nu}^I$	$Q_{\varphi d}$	$(\varphi^\dagger i \overleftrightarrow{D}_\mu \varphi)(\bar{d}_p \gamma^\mu d_r)$
$Q_{\varphi \tilde{W}B}$	$\varphi^\dagger \tau^I \varphi \tilde{W}_{\mu\nu}^I B^{\mu\nu}$	Q_{dB}	$(\bar{q}_p \sigma^{\mu\nu} d_r) \varphi B_{\mu\nu}$	$Q_{\varphi ud}$	$i(\tilde{\varphi}^\dagger D_\mu \varphi)(\bar{u}_p \gamma^\mu d_r)$




Bosonic operators

$\phi^n D^m$

[Grzadkowski et al.; JHEP 10 (2010) 085]

X^3		φ^6 and $\varphi^4 D^2$		$\psi^2 \varphi^3$	
Q_G	$f^{ABC} G_\mu^{A\nu} G_\nu^{B\rho} G_\rho^{C\mu}$	Q_φ	$(\varphi^\dagger \varphi)^3$	$Q_{e\varphi}$	$(\varphi^\dagger \varphi)(\bar{l}_p e_r \varphi)$
$Q_{\tilde{G}}$	$f^{ABC} \tilde{G}_\mu^{A\nu} G_\nu^{B\rho} G_\rho^{C\mu}$	$Q_{\varphi\Box}$	$(\varphi^\dagger \varphi)\Box(\varphi^\dagger \varphi)$	$Q_{u\varphi}$	$(\varphi^\dagger \varphi)(\bar{q}_p u_r \tilde{\varphi})$
Q_W	$\varepsilon^{IJK} W_\mu^{I\nu} W_\nu^{J\rho} W_\rho^{K\mu}$	$Q_{\varphi D}$	$(\varphi^\dagger D^\mu \varphi)^* (\varphi^\dagger D_\mu \varphi)$	$Q_{d\varphi}$	$(\varphi^\dagger \varphi)(\bar{q}_p d_r \varphi)$
$Q_{\tilde{W}}$	$\varepsilon^{IJK} \tilde{W}_\mu^{I\nu} W_\nu^{J\rho} W_\rho^{K\mu}$				
$X^2 \varphi^2$		$\psi^2 X \varphi$		$\psi^2 \varphi^2 D$	
$Q_{\varphi G}$	$\varphi^\dagger \varphi G_{\mu\nu}^A G^{A\mu\nu}$	Q_{eW}	$(\bar{l}_p \sigma^{\mu\nu} e_r) \tau^I \varphi W_{\mu\nu}^I$	$Q_{\varphi l}^{(1)}$	$(\varphi^\dagger i \overleftrightarrow{D}_\mu \varphi)(\bar{l}_p \gamma^\mu l_r)$
$Q_{\varphi \tilde{G}}$	$\varphi^\dagger \varphi \tilde{G}_{\mu\nu}^A G^{A\mu\nu}$	Q_{eB}	$(\bar{l}_p \sigma^{\mu\nu} e_r) \varphi B_{\mu\nu}$	$Q_{\varphi l}^{(3)}$	$(\varphi^\dagger i \overleftrightarrow{D}_\mu^I \varphi)(\bar{l}_p \tau^I \gamma^\mu l_r)$
$Q_{\varphi W}$	$\varphi^\dagger \varphi W_{\mu\nu}^I W^{I\mu\nu}$	Q_{uG}	$(\bar{q}_p \sigma^{\mu\nu} T^A u_r) \tilde{\varphi} G_{\mu\nu}^A$	$Q_{\varphi e}$	$(\varphi^\dagger i \overleftrightarrow{D}_\mu \varphi)(\bar{e}_p \gamma^\mu e_r)$
$Q_{\varphi \tilde{W}}$	$\varphi^\dagger \varphi \tilde{W}_{\mu\nu}^I W^{I\mu\nu}$	Q_{uW}	$(\bar{q}_p \sigma^{\mu\nu} u_r) \tau^I \tilde{\varphi} W_{\mu\nu}^I$	$Q_{\varphi q}^{(1)}$	$(\varphi^\dagger i \overleftrightarrow{D}_\mu \varphi)(\bar{q}_p \gamma^\mu q_r)$
$Q_{\varphi B}$	$\varphi^\dagger \varphi B_{\mu\nu} B^{\mu\nu}$	Q_{uB}	$(\bar{q}_p \sigma^{\mu\nu} u_r) \tilde{\varphi} B_{\mu\nu}$	$Q_{\varphi q}^{(3)}$	$(\varphi^\dagger i \overleftrightarrow{D}_\mu^I \varphi)(\bar{q}_p \tau^I \gamma^\mu q_r)$
$Q_{\varphi \tilde{B}}$	$\varphi^\dagger \varphi \tilde{B}_{\mu\nu} B^{\mu\nu}$	Q_{dG}	$(\bar{q}_p \sigma^{\mu\nu} T^A d_r) \varphi G_{\mu\nu}^A$	$Q_{\varphi u}$	$(\varphi^\dagger i \overleftrightarrow{D}_\mu \varphi)(\bar{u}_p \gamma^\mu u_r)$
$Q_{\varphi WB}$	$\varphi^\dagger \tau^I \varphi W_{\mu\nu}^I B^{\mu\nu}$	Q_{dW}	$(\bar{q}_p \sigma^{\mu\nu} d_r) \tau^I \varphi W_{\mu\nu}^I$	$Q_{\varphi d}$	$(\varphi^\dagger i \overleftrightarrow{D}_\mu \varphi)(\bar{d}_p \gamma^\mu d_r)$
$Q_{\varphi \tilde{W}B}$	$\varphi^\dagger \tau^I \varphi \tilde{W}_{\mu\nu}^I B^{\mu\nu}$	Q_{dB}	$(\bar{q}_p \sigma^{\mu\nu} d_r) \varphi B_{\mu\nu}$	$Q_{\varphi ud}$	$i(\tilde{\varphi}^\dagger D_\mu \varphi)(\bar{u}_p \gamma^\mu d_r)$




$\phi^2 X_{\mu\nu}^2$

Bosonic operators


$\phi^n D^m$

[Grzadkowski et al.; JHEP 10 (2010) 085]

X^3		φ^6 and $\varphi^4 D^2$		$\psi^2 \varphi^3$	
Q_G	$f^{ABC} G_{\mu}^{A\nu} G_{\nu}^{B\rho} G_{\rho}^{C\mu}$	Q_{φ}	$(\varphi^\dagger \varphi)^3$	$Q_{e\varphi}$	$(\varphi^\dagger \varphi)(\bar{l}_p e_r \varphi)$
$Q_{\tilde{G}}$	$f^{ABC} \tilde{G}_{\mu}^{A\nu} G_{\nu}^{B\rho} G_{\rho}^{C\mu}$	$Q_{\varphi\Box}$	$(\varphi^\dagger \varphi)\Box(\varphi^\dagger \varphi)$	$Q_{u\varphi}$	$(\varphi^\dagger \varphi)(\bar{q}_p u_r \tilde{\varphi})$
Q_W	$\varepsilon^{IJK} W_{\mu}^{I\nu} W_{\nu}^{J\rho} W_{\rho}^{K\mu}$	$Q_{\varphi D}$	$(\varphi^\dagger D^\mu \varphi)^* (\varphi^\dagger D_\mu \varphi)$	$Q_{d\varphi}$	$(\varphi^\dagger \varphi)(\bar{q}_p d_r \varphi)$
$Q_{\tilde{W}}$	$\varepsilon^{IJK} \tilde{W}_{\mu}^{I\nu} W_{\nu}^{J\rho} W_{\rho}^{K\mu}$				
$X^2 \varphi^2$		$\psi^2 X \varphi$		$\psi^2 \varphi^2 D$	
$Q_{\varphi G}$	$\varphi^\dagger \varphi G_{\mu\nu}^A G^{A\mu\nu}$	Q_{eW}	$(\bar{l}_p \sigma^{\mu\nu} e_r) \tau^I \varphi W_{\mu\nu}^I$	$Q_{\varphi l}^{(1)}$	$(\varphi^\dagger i \overleftrightarrow{D}_\mu \varphi)(\bar{l}_p \gamma^\mu l_r)$
$Q_{\varphi \tilde{G}}$	$\varphi^\dagger \varphi \tilde{G}_{\mu\nu}^A G^{A\mu\nu}$	Q_{eB}	$(\bar{l}_p \sigma^{\mu\nu} e_r) \varphi B_{\mu\nu}$	$Q_{\varphi l}^{(3)}$	$(\varphi^\dagger i \overleftrightarrow{D}_\mu^I \varphi)(\bar{l}_p \tau^I \gamma^\mu l_r)$
$Q_{\varphi W}$	$\varphi^\dagger \varphi W_{\mu\nu}^I W^{I\mu\nu}$	Q_{uG}	$(\bar{q}_p \sigma^{\mu\nu} T^A u_r) \tilde{\varphi} G_{\mu\nu}^A$	$Q_{\varphi e}$	$(\varphi^\dagger i \overleftrightarrow{D}_\mu \varphi)(\bar{e}_p \gamma^\mu e_r)$
$Q_{\varphi \tilde{W}}$	$\varphi^\dagger \varphi \tilde{W}_{\mu\nu}^I W^{I\mu\nu}$	Q_{uW}	$(\bar{q}_p \sigma^{\mu\nu} u_r) \tau^I \tilde{\varphi} W_{\mu\nu}^I$	$Q_{\varphi q}^{(1)}$	$(\varphi^\dagger i \overleftrightarrow{D}_\mu \varphi)(\bar{q}_p \gamma^\mu q_r)$
$Q_{\varphi B}$	$\varphi^\dagger \varphi B_{\mu\nu} B^{\mu\nu}$	Q_{uB}	$(\bar{q}_p \sigma^{\mu\nu} u_r) \tilde{\varphi} B_{\mu\nu}$	$Q_{\varphi q}^{(3)}$	$(\varphi^\dagger i \overleftrightarrow{D}_\mu^I \varphi)(\bar{q}_p \tau^I \gamma^\mu q_r)$
$Q_{\varphi \tilde{B}}$	$\varphi^\dagger \varphi \tilde{B}_{\mu\nu} B^{\mu\nu}$	Q_{dG}	$(\bar{q}_p \sigma^{\mu\nu} T^A d_r) \varphi G_{\mu\nu}^A$	$Q_{\varphi u}$	$(\varphi^\dagger i \overleftrightarrow{D}_\mu \varphi)(\bar{u}_p \gamma^\mu u_r)$
$Q_{\varphi WB}$	$\varphi^\dagger \tau^I \varphi W_{\mu\nu}^I B^{\mu\nu}$	Q_{dW}	$(\bar{q}_p \sigma^{\mu\nu} d_r) \tau^I \varphi W_{\mu\nu}^I$	$Q_{\varphi d}$	$(\varphi^\dagger i \overleftrightarrow{D}_\mu \varphi)(\bar{d}_p \gamma^\mu d_r)$
$Q_{\varphi \tilde{W}B}$	$\varphi^\dagger \tau^I \varphi \tilde{W}_{\mu\nu}^I B^{\mu\nu}$	Q_{dB}	$(\bar{q}_p \sigma^{\mu\nu} d_r) \varphi B_{\mu\nu}$	$Q_{\varphi ud}$	$i(\tilde{\varphi}^\dagger D_\mu \varphi)(\bar{u}_p \gamma^\mu d_r)$



$\phi^2 X_{\mu\nu}^2$





EW precision tests

Bosonic operators


$\phi^n D^m$

[Grzadkowski et al.; JHEP 10 (2010) 085]


X^3		φ^6 and $\varphi^4 D^2$		$\psi^2 \varphi^3$	
Q_G	$f^{ABC} G_\mu^{A\nu} G_\nu^{B\rho} G_\rho^{C\mu}$	Q_φ	$(\varphi^\dagger \varphi)^3$	$Q_{e\varphi}$	$(\varphi^\dagger \varphi)(\bar{l}_p e_r \varphi)$
$Q_{\tilde{G}}$	$f^{ABC} \tilde{G}_\mu^{A\nu} G_\nu^{B\rho} G_\rho^{C\mu}$	$Q_{\varphi\Box}$	$(\varphi^\dagger \varphi)\Box(\varphi^\dagger \varphi)$	$Q_{u\varphi}$	$(\varphi^\dagger \varphi)(\bar{q}_p u_r \tilde{\varphi})$
Q_W	$\varepsilon^{IJK} W_\mu^{I\nu} W_\nu^{J\rho} W_\rho^{K\mu}$	$Q_{\varphi D}$	$(\varphi^\dagger D^\mu \varphi)^* (\varphi^\dagger D_\mu \varphi)$	$Q_{d\varphi}$	$(\varphi^\dagger \varphi)(\bar{q}_p d_r \varphi)$
$Q_{\tilde{W}}$	$\varepsilon^{IJK} \tilde{W}_\mu^{I\nu} W_\nu^{J\rho} W_\rho^{K\mu}$				
$X^2 \varphi^2$		$\psi^2 X \varphi$		$\psi^2 \varphi^2 D$	
$Q_{\varphi G}$	$\varphi^\dagger \varphi G_{\mu\nu}^A G^{A\mu\nu}$	Q_{eW}	$(\bar{l}_p \sigma^{\mu\nu} e_r) \tau^I \varphi W_{\mu\nu}^I$	$Q_{\varphi l}^{(1)}$	$(\varphi^\dagger i \overleftrightarrow{D}_\mu \varphi)(\bar{l}_p \gamma^\mu l_r)$
$Q_{\varphi \tilde{G}}$	$\varphi^\dagger \varphi \tilde{G}_{\mu\nu}^A G^{A\mu\nu}$	Q_{eB}	$(\bar{l}_p \sigma^{\mu\nu} e_r) \varphi B_{\mu\nu}$	$Q_{\varphi l}^{(3)}$	$(\varphi^\dagger i \overleftrightarrow{D}_\mu^I \varphi)(\bar{l}_p \tau^I \gamma^\mu l_r)$
$Q_{\varphi W}$	$\varphi^\dagger \varphi W_{\mu\nu}^I W^{I\mu\nu}$	Q_{uG}	$(\bar{q}_p \sigma^{\mu\nu} T^A u_r) \tilde{\varphi} G_{\mu\nu}^A$	$Q_{\varphi e}$	$(\varphi^\dagger i \overleftrightarrow{D}_\mu \varphi)(\bar{e}_p \gamma^\mu e_r)$
$Q_{\varphi \tilde{W}}$	$\varphi^\dagger \varphi \tilde{W}_{\mu\nu}^I W^{I\mu\nu}$	Q_{uW}	$(\bar{q}_p \sigma^{\mu\nu} u_r) \tau^I \tilde{\varphi} W_{\mu\nu}^I$	$Q_{\varphi q}^{(1)}$	$(\varphi^\dagger i \overleftrightarrow{D}_\mu \varphi)(\bar{q}_p \gamma^\mu q_r)$
$Q_{\varphi B}$	$\varphi^\dagger \varphi B_{\mu\nu} B^{\mu\nu}$	Q_{uB}	$(\bar{q}_p \sigma^{\mu\nu} u_r) \tilde{\varphi} B_{\mu\nu}$	$Q_{\varphi q}^{(3)}$	$(\varphi^\dagger i \overleftrightarrow{D}_\mu^I \varphi)(\bar{q}_p \tau^I \gamma^\mu q_r)$
$Q_{\varphi \tilde{B}}$	$\varphi^\dagger \varphi \tilde{B}_{\mu\nu} B^{\mu\nu}$	Q_{dG}	$(\bar{q}_p \sigma^{\mu\nu} T^A d_r) \varphi G_{\mu\nu}^A$	$Q_{\varphi u}$	$(\varphi^\dagger i \overleftrightarrow{D}_\mu \varphi)(\bar{u}_p \gamma^\mu u_r)$
$Q_{\varphi WB}$	$\varphi^\dagger \tau^I \varphi W_{\mu\nu}^I B^{\mu\nu}$	Q_{dW}	$(\bar{q}_p \sigma^{\mu\nu} d_r) \tau^I \varphi W_{\mu\nu}^I$	$Q_{\varphi d}$	$(\varphi^\dagger i \overleftrightarrow{D}_\mu \varphi)(\bar{d}_p \gamma^\mu d_r)$
$Q_{\varphi \tilde{W}B}$	$\varphi^\dagger \tau^I \varphi \tilde{W}_{\mu\nu}^I B^{\mu\nu}$	Q_{dB}	$(\bar{q}_p \sigma^{\mu\nu} d_r) \varphi B_{\mu\nu}$	$Q_{\varphi ud}$	$i(\tilde{\varphi}^\dagger D_\mu \varphi)(\bar{u}_p \gamma^\mu d_r)$



$\phi^2 X_{\mu\nu}^2$



EW precision tests






EW Higgs production & decay

Bosonic operators


$\phi^n D^m$

[Grzadkowski et al.; JHEP 10 (2010) 085]



X^3		φ^6 and $\varphi^4 D^2$		$\psi^2 \varphi^3$	
Q_G	$f^{ABC} G_\mu^{A\nu} G_\nu^{B\rho} G_\rho^{C\mu}$	Q_φ	$(\varphi^\dagger \varphi)^3$	$Q_{e\varphi}$	$(\varphi^\dagger \varphi)(\bar{l}_p e_r \varphi)$
$Q_{\tilde{G}}$	$f^{ABC} \tilde{G}_\mu^{A\nu} G_\nu^{B\rho} G_\rho^{C\mu}$	$Q_{\varphi\Box}$	$(\varphi^\dagger \varphi)\Box(\varphi^\dagger \varphi)$	$Q_{u\varphi}$	$(\varphi^\dagger \varphi)(\bar{q}_p u_r \tilde{\varphi})$
Q_W	$\varepsilon^{IJK} W_\mu^{I\nu} W_\nu^{J\rho} W_\rho^{K\mu}$	$Q_{\varphi D}$	$(\varphi^\dagger D^\mu \varphi)^* (\varphi^\dagger D_\mu \varphi)$	$Q_{d\varphi}$	$(\varphi^\dagger \varphi)(\bar{q}_p d_r \varphi)$
$Q_{\tilde{W}}$	$\varepsilon^{IJK} \tilde{W}_\mu^{I\nu} W_\nu^{J\rho} W_\rho^{K\mu}$				
$X^2 \varphi^2$		$\psi^2 X \varphi$		$\psi^2 \varphi^2 D$	
$Q_{\varphi G}$	$\varphi^\dagger \varphi G_{\mu\nu}^A G^{A\mu\nu}$	Q_{eW}	$(\bar{l}_p \sigma^{\mu\nu} e_r) \tau^I \varphi W_{\mu\nu}^I$	$Q_{\varphi l}^{(1)}$	$(\varphi^\dagger i \overleftrightarrow{D}_\mu \varphi)(\bar{l}_p \gamma^\mu l_r)$
$Q_{\varphi \tilde{G}}$	$\varphi^\dagger \varphi \tilde{G}_{\mu\nu}^A G^{A\mu\nu}$	Q_{eB}	$(\bar{l}_p \sigma^{\mu\nu} e_r) \varphi B_{\mu\nu}$	$Q_{\varphi l}^{(3)}$	$(\varphi^\dagger i \overleftrightarrow{D}_\mu^I \varphi)(\bar{l}_p \tau^I \gamma^\mu l_r)$
$Q_{\varphi W}$	$\varphi^\dagger \varphi W_{\mu\nu}^I W^{I\mu\nu}$	Q_{uG}	$(\bar{q}_p \sigma^{\mu\nu} T^A u_r) \tilde{\varphi} G_{\mu\nu}^A$	$Q_{\varphi e}$	$(\varphi^\dagger i \overleftrightarrow{D}_\mu \varphi)(\bar{e}_p \gamma^\mu e_r)$
$Q_{\varphi \tilde{W}}$	$\varphi^\dagger \varphi \tilde{W}_{\mu\nu}^I W^{I\mu\nu}$	Q_{uW}	$(\bar{q}_p \sigma^{\mu\nu} u_r) \tau^I \tilde{\varphi} W_{\mu\nu}^I$	$Q_{\varphi q}^{(1)}$	$(\varphi^\dagger i \overleftrightarrow{D}_\mu \varphi)(\bar{q}_p \gamma^\mu q_r)$
$Q_{\varphi B}$	$\varphi^\dagger \varphi B_{\mu\nu} B^{\mu\nu}$	Q_{uB}	$(\bar{q}_p \sigma^{\mu\nu} u_r) \tilde{\varphi} B_{\mu\nu}$	$Q_{\varphi q}^{(3)}$	$(\varphi^\dagger i \overleftrightarrow{D}_\mu^I \varphi)(\bar{q}_p \tau^I \gamma^\mu q_r)$
$Q_{\varphi \tilde{B}}$	$\varphi^\dagger \varphi \tilde{B}_{\mu\nu} B^{\mu\nu}$	Q_{dG}	$(\bar{q}_p \sigma^{\mu\nu} T^A d_r) \varphi G_{\mu\nu}^A$	$Q_{\varphi u}$	$(\varphi^\dagger i \overleftrightarrow{D}_\mu \varphi)(\bar{u}_p \gamma^\mu u_r)$
$Q_{\varphi WB}$	$\varphi^\dagger \tau^I \varphi W_{\mu\nu}^I B^{\mu\nu}$	Q_{dW}	$(\bar{q}_p \sigma^{\mu\nu} d_r) \tau^I \varphi W_{\mu\nu}^I$	$Q_{\varphi d}$	$(\varphi^\dagger i \overleftrightarrow{D}_\mu \varphi)(\bar{d}_p \gamma^\mu d_r)$
$Q_{\varphi \tilde{W}B}$	$\varphi^\dagger \tau^I \varphi \tilde{W}_{\mu\nu}^I B^{\mu\nu}$	Q_{dB}	$(\bar{q}_p \sigma^{\mu\nu} d_r) \varphi B_{\mu\nu}$	$Q_{\varphi ud}$	$i(\tilde{\varphi}^\dagger D_\mu \varphi)(\bar{u}_p \gamma^\mu d_r)$




EW precision tests



EW Higgs production & decay



Gluon fusion: Higgs, di-Higgs





$\phi^2 X_{\mu\nu}^2$

Yukawa/dipole operators

$$\phi^3 \bar{\psi}_L \psi_R$$

[Grzadkowski et al.; JHEP 10 (2010) 085]

X^3		φ^6 and $\varphi^4 D^2$		$\psi^2 \varphi^3$	
Q_G	$f^{ABC} G_{\mu}^{A\nu} G_{\nu}^{B\rho} G_{\rho}^{C\mu}$	Q_{φ}	$(\varphi^\dagger \varphi)^3$	$Q_{e\varphi}$	$(\varphi^\dagger \varphi)(\bar{l}_p e_r \varphi)$
$Q_{\tilde{G}}$	$f^{ABC} \tilde{G}_{\mu}^{A\nu} G_{\nu}^{B\rho} G_{\rho}^{C\mu}$	$Q_{\varphi\Box}$	$(\varphi^\dagger \varphi)\Box(\varphi^\dagger \varphi)$	$Q_{u\varphi}$	$(\varphi^\dagger \varphi)(\bar{q}_p u_r \tilde{\varphi})$
Q_W	$\varepsilon^{IJK} W_{\mu}^{I\nu} W_{\nu}^{J\rho} W_{\rho}^{K\mu}$	$Q_{\varphi D}$	$(\varphi^\dagger D^\mu \varphi)^* (\varphi^\dagger D_\mu \varphi)$	$Q_{d\varphi}$	$(\varphi^\dagger \varphi)(\bar{q}_p d_r \varphi)$
$Q_{\tilde{W}}$	$\varepsilon^{IJK} \tilde{W}_{\mu}^{I\nu} W_{\nu}^{J\rho} W_{\rho}^{K\mu}$				
$X^2 \varphi^2$		$\psi^2 X \varphi$		$\psi^2 \varphi^2 D$	
$Q_{\varphi G}$	$\varphi^\dagger \varphi G_{\mu\nu}^A G^{A\mu\nu}$	Q_{eW}	$(\bar{l}_p \sigma^{\mu\nu} e_r) \tau^I \varphi W_{\mu\nu}^I$	$Q_{\varphi l}^{(1)}$	$(\varphi^\dagger i \overleftrightarrow{D}_\mu \varphi)(\bar{l}_p \gamma^\mu l_r)$
$Q_{\varphi \tilde{G}}$	$\varphi^\dagger \varphi \tilde{G}_{\mu\nu}^A G^{A\mu\nu}$	Q_{eB}	$(\bar{l}_p \sigma^{\mu\nu} e_r) \varphi B_{\mu\nu}$	$Q_{\varphi l}^{(3)}$	$(\varphi^\dagger i \overleftrightarrow{D}_\mu^I \varphi)(\bar{l}_p \tau^I \gamma^\mu l_r)$
$Q_{\varphi W}$	$\varphi^\dagger \varphi W_{\mu\nu}^I W^{I\mu\nu}$	Q_{uG}	$(\bar{q}_p \sigma^{\mu\nu} T^A u_r) \tilde{\varphi} G_{\mu\nu}^A$	$Q_{\varphi e}$	$(\varphi^\dagger i \overleftrightarrow{D}_\mu \varphi)(\bar{e}_p \gamma^\mu e_r)$
$Q_{\varphi \tilde{W}}$	$\varphi^\dagger \varphi \tilde{W}_{\mu\nu}^I W^{I\mu\nu}$	Q_{uW}	$(\bar{q}_p \sigma^{\mu\nu} u_r) \tau^I \tilde{\varphi} W_{\mu\nu}^I$	$Q_{\varphi q}^{(1)}$	$(\varphi^\dagger i \overleftrightarrow{D}_\mu \varphi)(\bar{q}_p \gamma^\mu q_r)$
$Q_{\varphi B}$	$\varphi^\dagger \varphi B_{\mu\nu} B^{\mu\nu}$	Q_{uB}	$(\bar{q}_p \sigma^{\mu\nu} u_r) \tilde{\varphi} B_{\mu\nu}$	$Q_{\varphi q}^{(3)}$	$(\varphi^\dagger i \overleftrightarrow{D}_\mu^I \varphi)(\bar{q}_p \tau^I \gamma^\mu q_r)$
$Q_{\varphi \tilde{B}}$	$\varphi^\dagger \varphi \tilde{B}_{\mu\nu} B^{\mu\nu}$	Q_{dG}	$(\bar{q}_p \sigma^{\mu\nu} T^A d_r) \varphi G_{\mu\nu}^A$	$Q_{\varphi u}$	$(\varphi^\dagger i \overleftrightarrow{D}_\mu \varphi)(\bar{u}_p \gamma^\mu u_r)$
$Q_{\varphi WB}$	$\varphi^\dagger \tau^I \varphi W_{\mu\nu}^I B^{\mu\nu}$	Q_{dW}	$(\bar{q}_p \sigma^{\mu\nu} d_r) \tau^I \varphi W_{\mu\nu}^I$	$Q_{\varphi d}$	$(\varphi^\dagger i \overleftrightarrow{D}_\mu \varphi)(\bar{d}_p \gamma^\mu d_r)$
$Q_{\varphi \tilde{W}B}$	$\varphi^\dagger \tau^I \varphi \tilde{W}_{\mu\nu}^I B^{\mu\nu}$	Q_{dB}	$(\bar{q}_p \sigma^{\mu\nu} d_r) \varphi B_{\mu\nu}$	$Q_{\varphi ud}$	$i(\tilde{\varphi}^\dagger D_\mu \varphi)(\bar{u}_p \gamma^\mu d_r)$





Yukawa/dipole operators

[Grzadkowski et al.; JHEP 10 (2010) 085]


X^3		φ^6 and $\varphi^4 D^2$		$\psi^2 \varphi^3$	
Q_G	$f^{ABC} G_{\mu}^{A\nu} G_{\nu}^{B\rho} G_{\rho}^{C\mu}$	Q_{φ}	$(\varphi^\dagger \varphi)^3$	$Q_{e\varphi}$	$(\varphi^\dagger \varphi)(\bar{l}_p e_r \varphi)$
$Q_{\tilde{G}}$	$f^{ABC} \tilde{G}_{\mu}^{A\nu} G_{\nu}^{B\rho} G_{\rho}^{C\mu}$	$Q_{\varphi\Box}$	$(\varphi^\dagger \varphi)\Box(\varphi^\dagger \varphi)$	$Q_{u\varphi}$	$(\varphi^\dagger \varphi)(\bar{q}_p u_r \tilde{\varphi})$
Q_W	$\varepsilon^{IJK} W_{\mu}^{I\nu} W_{\nu}^{J\rho} W_{\rho}^{K\mu}$	$Q_{\varphi D}$	$(\varphi^\dagger D^\mu \varphi)^* (\varphi^\dagger D_\mu \varphi)$	$Q_{d\varphi}$	$(\varphi^\dagger \varphi)(\bar{q}_p d_r \varphi)$
$Q_{\tilde{W}}$	$\varepsilon^{IJK} \tilde{W}_{\mu}^{I\nu} W_{\nu}^{J\rho} W_{\rho}^{K\mu}$				
$X^2 \varphi^2$		$\psi^2 X \varphi$		$\psi^2 \varphi^2 D$	
$Q_{\varphi G}$	$\varphi^\dagger \varphi G_{\mu\nu}^A G^{A\mu\nu}$	Q_{eW}	$(\bar{l}_p \sigma^{\mu\nu} e_r) \tau^I \varphi W_{\mu\nu}^I$	$Q_{\varphi l}^{(1)}$	$(\varphi^\dagger i \overleftrightarrow{D}_\mu \varphi)(\bar{l}_p \gamma^\mu l_r)$
$Q_{\varphi \tilde{G}}$	$\varphi^\dagger \varphi \tilde{G}_{\mu\nu}^A G^{A\mu\nu}$	Q_{eB}	$(\bar{l}_p \sigma^{\mu\nu} e_r) \varphi B_{\mu\nu}$	$Q_{\varphi l}^{(3)}$	$(\varphi^\dagger i \overleftrightarrow{D}_\mu^I \varphi)(\bar{l}_p \tau^I \gamma^\mu l_r)$
$Q_{\varphi W}$	$\varphi^\dagger \varphi W_{\mu\nu}^I W^{I\mu\nu}$	Q_{uG}	$(\bar{q}_p \sigma^{\mu\nu} T^A u_r) \tilde{\varphi} G_{\mu\nu}^A$	$Q_{\varphi e}$	$(\varphi^\dagger i \overleftrightarrow{D}_\mu \varphi)(\bar{e}_p \gamma^\mu e_r)$
$Q_{\varphi \tilde{W}}$	$\varphi^\dagger \varphi \tilde{W}_{\mu\nu}^I W^{I\mu\nu}$	Q_{uW}	$(\bar{q}_p \sigma^{\mu\nu} u_r) \tau^I \tilde{\varphi} W_{\mu\nu}^I$	$Q_{\varphi q}^{(1)}$	$(\varphi^\dagger i \overleftrightarrow{D}_\mu \varphi)(\bar{q}_p \gamma^\mu q_r)$
$Q_{\varphi B}$	$\varphi^\dagger \varphi B_{\mu\nu} B^{\mu\nu}$	Q_{uB}	$(\bar{q}_p \sigma^{\mu\nu} u_r) \tilde{\varphi} B_{\mu\nu}$	$Q_{\varphi q}^{(3)}$	$(\varphi^\dagger i \overleftrightarrow{D}_\mu^I \varphi)(\bar{q}_p \tau^I \gamma^\mu q_r)$
$Q_{\varphi \tilde{B}}$	$\varphi^\dagger \varphi \tilde{B}_{\mu\nu} B^{\mu\nu}$	Q_{dG}	$(\bar{q}_p \sigma^{\mu\nu} T^A d_r) \varphi G_{\mu\nu}^A$	$Q_{\varphi u}$	$(\varphi^\dagger i \overleftrightarrow{D}_\mu \varphi)(\bar{u}_p \gamma^\mu u_r)$
$Q_{\varphi WB}$	$\varphi^\dagger \tau^I \varphi W_{\mu\nu}^I B^{\mu\nu}$	Q_{dW}	$(\bar{q}_p \sigma^{\mu\nu} d_r) \tau^I \varphi W_{\mu\nu}^I$	$Q_{\varphi d}$	$(\varphi^\dagger i \overleftrightarrow{D}_\mu \varphi)(\bar{d}_p \gamma^\mu d_r)$
$Q_{\varphi \tilde{W}B}$	$\varphi^\dagger \tau^I \varphi \tilde{W}_{\mu\nu}^I B^{\mu\nu}$	Q_{dB}	$(\bar{q}_p \sigma^{\mu\nu} d_r) \varphi B_{\mu\nu}$	$Q_{\varphi ud}$	$i(\tilde{\varphi}^\dagger D_\mu \varphi)(\bar{u}_p \gamma^\mu d_r)$

$$\phi^3 \bar{\psi}_L \psi_R$$



$$H \rightarrow f\bar{f}$$

$$\phi \bar{\psi}_L \sigma_{\mu\nu} \psi_R X_{\mu\nu}$$





Yukawa/dipole operators

[Grzadkowski et al.; JHEP 10 (2010) 085]


X^3		φ^6 and $\varphi^4 D^2$		$\psi^2 \varphi^3$	
Q_G	$f^{ABC} G_{\mu}^{A\nu} G_{\nu}^{B\rho} G_{\rho}^{C\mu}$	Q_{φ}	$(\varphi^\dagger \varphi)^3$	$Q_{e\varphi}$	$(\varphi^\dagger \varphi)(\bar{l}_p e_r \varphi)$
$Q_{\tilde{G}}$	$f^{ABC} \tilde{G}_{\mu}^{A\nu} G_{\nu}^{B\rho} G_{\rho}^{C\mu}$	$Q_{\varphi\Box}$	$(\varphi^\dagger \varphi)\Box(\varphi^\dagger \varphi)$	$Q_{u\varphi}$	$(\varphi^\dagger \varphi)(\bar{q}_p u_r \tilde{\varphi})$
Q_W	$\varepsilon^{IJK} W_{\mu}^{I\nu} W_{\nu}^{J\rho} W_{\rho}^{K\mu}$	$Q_{\varphi D}$	$(\varphi^\dagger D^\mu \varphi)^* (\varphi^\dagger D_\mu \varphi)$	$Q_{d\varphi}$	$(\varphi^\dagger \varphi)(\bar{q}_p d_r \varphi)$
$Q_{\tilde{W}}$	$\varepsilon^{IJK} \tilde{W}_{\mu}^{I\nu} W_{\nu}^{J\rho} W_{\rho}^{K\mu}$				
$X^2 \varphi^2$		$\psi^2 X \varphi$		$\psi^2 \varphi^2 D$	
$Q_{\varphi G}$	$\varphi^\dagger \varphi G_{\mu\nu}^A G^{A\mu\nu}$	Q_{eW}	$(\bar{l}_p \sigma^{\mu\nu} e_r) \tau^I \varphi W_{\mu\nu}^I$	$Q_{\varphi l}^{(1)}$	$(\varphi^\dagger i \overleftrightarrow{D}_\mu \varphi)(\bar{l}_p \gamma^\mu l_r)$
$Q_{\varphi \tilde{G}}$	$\varphi^\dagger \varphi \tilde{G}_{\mu\nu}^A G^{A\mu\nu}$	Q_{eB}	$(\bar{l}_p \sigma^{\mu\nu} e_r) \varphi B_{\mu\nu}$	$Q_{\varphi l}^{(3)}$	$(\varphi^\dagger i \overleftrightarrow{D}_\mu^I \varphi)(\bar{l}_p \tau^I \gamma^\mu l_r)$
$Q_{\varphi W}$	$\varphi^\dagger \varphi W_{\mu\nu}^I W^{I\mu\nu}$	Q_{uG}	$(\bar{q}_p \sigma^{\mu\nu} T^A u_r) \tilde{\varphi} G_{\mu\nu}^A$	$Q_{\varphi e}$	$(\varphi^\dagger i \overleftrightarrow{D}_\mu \varphi)(\bar{e}_p \gamma^\mu e_r)$
$Q_{\varphi \tilde{W}}$	$\varphi^\dagger \varphi \tilde{W}_{\mu\nu}^I W^{I\mu\nu}$	Q_{uW}	$(\bar{q}_p \sigma^{\mu\nu} u_r) \tau^I \tilde{\varphi} W_{\mu\nu}^I$	$Q_{\varphi q}^{(1)}$	$(\varphi^\dagger i \overleftrightarrow{D}_\mu \varphi)(\bar{q}_p \gamma^\mu q_r)$
$Q_{\varphi B}$	$\varphi^\dagger \varphi B_{\mu\nu} B^{\mu\nu}$	Q_{uB}	$(\bar{q}_p \sigma^{\mu\nu} u_r) \tilde{\varphi} B_{\mu\nu}$	$Q_{\varphi q}^{(3)}$	$(\varphi^\dagger i \overleftrightarrow{D}_\mu^I \varphi)(\bar{q}_p \tau^I \gamma^\mu q_r)$
$Q_{\varphi \tilde{B}}$	$\varphi^\dagger \varphi \tilde{B}_{\mu\nu} B^{\mu\nu}$	Q_{dG}	$(\bar{q}_p \sigma^{\mu\nu} T^A d_r) \varphi G_{\mu\nu}^A$	$Q_{\varphi u}$	$(\varphi^\dagger i \overleftrightarrow{D}_\mu \varphi)(\bar{u}_p \gamma^\mu u_r)$
$Q_{\varphi WB}$	$\varphi^\dagger \tau^I \varphi W_{\mu\nu}^I B^{\mu\nu}$	Q_{dW}	$(\bar{q}_p \sigma^{\mu\nu} d_r) \tau^I \varphi W_{\mu\nu}^I$	$Q_{\varphi d}$	$(\varphi^\dagger i \overleftrightarrow{D}_\mu \varphi)(\bar{d}_p \gamma^\mu d_r)$
$Q_{\varphi \tilde{W}B}$	$\varphi^\dagger \tau^I \varphi \tilde{W}_{\mu\nu}^I B^{\mu\nu}$	Q_{dB}	$(\bar{q}_p \sigma^{\mu\nu} d_r) \varphi B_{\mu\nu}$	$Q_{\varphi ud}$	$i(\tilde{\varphi}^\dagger D_\mu \varphi)(\bar{u}_p \gamma^\mu d_r)$

$$\phi^3 \bar{\psi}_L \psi_R$$



$$H \rightarrow f\bar{f}$$

$$\phi \bar{\psi}_L \sigma_{\mu\nu} \psi_R X_{\mu\nu}$$



$$Z, W, \gamma, g$$

Chiral flip: $f_L f_R$


⇒ Leading effects $\propto m_f$

⇒ 3rd generation couplings

Yukawa/dipole operators

[Grzadkowski et al.; JHEP 10 (2010) 085]


X^3		φ^6 and $\varphi^4 D^2$		$\psi^2 \varphi^3$	
Q_G	$f^{ABC} G_{\mu}^{A\nu} G_{\nu}^{B\rho} G_{\rho}^{C\mu}$	Q_{φ}	$(\varphi^\dagger \varphi)^3$	$Q_{e\varphi}$	$(\varphi^\dagger \varphi)(\bar{l}_p e_r \varphi)$
$Q_{\tilde{G}}$	$f^{ABC} \tilde{G}_{\mu}^{A\nu} G_{\nu}^{B\rho} G_{\rho}^{C\mu}$	$Q_{\varphi\Box}$	$(\varphi^\dagger \varphi)\Box(\varphi^\dagger \varphi)$	$Q_{u\varphi}$	$(\varphi^\dagger \varphi)(\bar{q}_p u_r \tilde{\varphi})$
Q_W	$\varepsilon^{IJK} W_{\mu}^{I\nu} W_{\nu}^{J\rho} W_{\rho}^{K\mu}$	$Q_{\varphi D}$	$(\varphi^\dagger D^\mu \varphi)^* (\varphi^\dagger D_\mu \varphi)$	$Q_{d\varphi}$	$(\varphi^\dagger \varphi)(\bar{q}_p d_r \varphi)$
$Q_{\tilde{W}}$	$\varepsilon^{IJK} \tilde{W}_{\mu}^{I\nu} W_{\nu}^{J\rho} W_{\rho}^{K\mu}$				
$X^2 \varphi^2$		$\psi^2 X \varphi$		$\psi^2 \varphi^2 D$	
$Q_{\varphi G}$	$\varphi^\dagger \varphi G_{\mu\nu}^A G^{A\mu\nu}$	Q_{eW}	$(\bar{l}_p \sigma^{\mu\nu} e_r) \tau^I \varphi W_{\mu\nu}^I$	$Q_{\varphi l}^{(1)}$	$(\varphi^\dagger i \overleftrightarrow{D}_\mu \varphi)(\bar{l}_p \gamma^\mu l_r)$
$Q_{\varphi \tilde{G}}$	$\varphi^\dagger \varphi \tilde{G}_{\mu\nu}^A G^{A\mu\nu}$	Q_{eB}	$(\bar{l}_p \sigma^{\mu\nu} e_r) \varphi B_{\mu\nu}$	$Q_{\varphi l}^{(3)}$	$(\varphi^\dagger i \overleftrightarrow{D}_\mu^I \varphi)(\bar{l}_p \tau^I \gamma^\mu l_r)$
$Q_{\varphi W}$	$\varphi^\dagger \varphi W_{\mu\nu}^I W^{I\mu\nu}$	Q_{uG}	$(\bar{q}_p \sigma^{\mu\nu} T^A u_r) \tilde{\varphi} G_{\mu\nu}^A$	$Q_{\varphi e}$	$(\varphi^\dagger i \overleftrightarrow{D}_\mu \varphi)(\bar{e}_p \gamma^\mu e_r)$
$Q_{\varphi \tilde{W}}$	$\varphi^\dagger \varphi \tilde{W}_{\mu\nu}^I W^{I\mu\nu}$	Q_{uW}	$(\bar{q}_p \sigma^{\mu\nu} u_r) \tau^I \tilde{\varphi} W_{\mu\nu}^I$	$Q_{\varphi q}^{(1)}$	$(\varphi^\dagger i \overleftrightarrow{D}_\mu \varphi)(\bar{q}_p \gamma^\mu q_r)$
$Q_{\varphi B}$	$\varphi^\dagger \varphi B_{\mu\nu} B^{\mu\nu}$	Q_{uB}	$(\bar{q}_p \sigma^{\mu\nu} u_r) \tilde{\varphi} B_{\mu\nu}$	$Q_{\varphi q}^{(3)}$	$(\varphi^\dagger i \overleftrightarrow{D}_\mu^I \varphi)(\bar{q}_p \tau^I \gamma^\mu q_r)$
$Q_{\varphi \tilde{B}}$	$\varphi^\dagger \varphi \tilde{B}_{\mu\nu} B^{\mu\nu}$	Q_{dG}	$(\bar{q}_p \sigma^{\mu\nu} T^A d_r) \varphi G_{\mu\nu}^A$	$Q_{\varphi u}$	$(\varphi^\dagger i \overleftrightarrow{D}_\mu \varphi)(\bar{u}_p \gamma^\mu u_r)$
$Q_{\varphi WB}$	$\varphi^\dagger \tau^I \varphi W_{\mu\nu}^I B^{\mu\nu}$	Q_{dW}	$(\bar{q}_p \sigma^{\mu\nu} d_r) \tau^I \varphi W_{\mu\nu}^I$	$Q_{\varphi d}$	$(\varphi^\dagger i \overleftrightarrow{D}_\mu \varphi)(\bar{d}_p \gamma^\mu d_r)$
$Q_{\varphi \tilde{W}B}$	$\varphi^\dagger \tau^I \varphi \tilde{W}_{\mu\nu}^I B^{\mu\nu}$	Q_{dB}	$(\bar{q}_p \sigma^{\mu\nu} d_r) \varphi B_{\mu\nu}$	$Q_{\varphi ud}$	$i(\tilde{\varphi}^\dagger D_\mu \varphi)(\bar{u}_p \gamma^\mu d_r)$




Chiral flip: $f_L f_R$

⇒ Leading effects $\propto m_f$


⇒ 3rd generation couplings



ggH



ggHH




ttH

Current operators

[Grzadkowski et al.; JHEP 10 (2010) 085]

X^3		φ^6 and $\varphi^4 D^2$		$\psi^2 \varphi^3$	
Q_G	$f^{ABC} G_{\mu}^{A\nu} G_{\nu}^{B\rho} G_{\rho}^{C\mu}$	Q_{φ}	$(\varphi^\dagger \varphi)^3$	$Q_{e\varphi}$	$(\varphi^\dagger \varphi)(\bar{l}_p e_r \varphi)$
$Q_{\tilde{G}}$	$f^{ABC} \tilde{G}_{\mu}^{A\nu} G_{\nu}^{B\rho} G_{\rho}^{C\mu}$	$Q_{\varphi\Box}$	$(\varphi^\dagger \varphi)\Box(\varphi^\dagger \varphi)$	$Q_{u\varphi}$	$(\varphi^\dagger \varphi)(\bar{q}_p u_r \tilde{\varphi})$
Q_W	$\varepsilon^{IJK} W_{\mu}^{I\nu} W_{\nu}^{J\rho} W_{\rho}^{K\mu}$	$Q_{\varphi D}$	$(\varphi^\dagger D^\mu \varphi)^* (\varphi^\dagger D_\mu \varphi)$	$Q_{d\varphi}$	$(\varphi^\dagger \varphi)(\bar{q}_p d_r \varphi)$
$Q_{\tilde{W}}$	$\varepsilon^{IJK} \tilde{W}_{\mu}^{I\nu} W_{\nu}^{J\rho} W_{\rho}^{K\mu}$				
$X^2 \varphi^2$		$\psi^2 X \varphi$		$\psi^2 \varphi^2 D$	
$Q_{\varphi G}$	$\varphi^\dagger \varphi G_{\mu\nu}^A G^{A\mu\nu}$	Q_{eW}	$(\bar{l}_p \sigma^{\mu\nu} e_r) \tau^I \varphi W_{\mu\nu}^I$	$Q_{\varphi l}^{(1)}$	$(\varphi^\dagger i \overleftrightarrow{D}_\mu \varphi)(\bar{l}_p \gamma^\mu l_r)$
$Q_{\varphi \tilde{G}}$	$\varphi^\dagger \varphi \tilde{G}_{\mu\nu}^A G^{A\mu\nu}$	Q_{eB}	$(\bar{l}_p \sigma^{\mu\nu} e_r) \varphi B_{\mu\nu}$	$Q_{\varphi l}^{(3)}$	$(\varphi^\dagger i \overleftrightarrow{D}_\mu^I \varphi)(\bar{l}_p \tau^I \gamma^\mu l_r)$
$Q_{\varphi W}$	$\varphi^\dagger \varphi W_{\mu\nu}^I W^{I\mu\nu}$	Q_{uG}	$(\bar{q}_p \sigma^{\mu\nu} T^A u_r) \tilde{\varphi} G_{\mu\nu}^A$	$Q_{\varphi e}$	$(\varphi^\dagger i \overleftrightarrow{D}_\mu \varphi)(\bar{e}_p \gamma^\mu e_r)$
$Q_{\varphi \tilde{W}}$	$\varphi^\dagger \varphi \tilde{W}_{\mu\nu}^I W^{I\mu\nu}$	Q_{uW}	$(\bar{q}_p \sigma^{\mu\nu} u_r) \tau^I \tilde{\varphi} W_{\mu\nu}^I$	$Q_{\varphi q}^{(1)}$	$(\varphi^\dagger i \overleftrightarrow{D}_\mu \varphi)(\bar{q}_p \gamma^\mu q_r)$
$Q_{\varphi B}$	$\varphi^\dagger \varphi B_{\mu\nu} B^{\mu\nu}$	Q_{uB}	$(\bar{q}_p \sigma^{\mu\nu} u_r) \tilde{\varphi} B_{\mu\nu}$	$Q_{\varphi q}^{(3)}$	$(\varphi^\dagger i \overleftrightarrow{D}_\mu^I \varphi)(\bar{q}_p \tau^I \gamma^\mu q_r)$
$Q_{\varphi \tilde{B}}$	$\varphi^\dagger \varphi \tilde{B}_{\mu\nu} B^{\mu\nu}$	Q_{dG}	$(\bar{q}_p \sigma^{\mu\nu} T^A d_r) \varphi G_{\mu\nu}^A$	$Q_{\varphi u}$	$(\varphi^\dagger i \overleftrightarrow{D}_\mu \varphi)(\bar{u}_p \gamma^\mu u_r)$
$Q_{\varphi WB}$	$\varphi^\dagger \tau^I \varphi W_{\mu\nu}^I B^{\mu\nu}$	Q_{dW}	$(\bar{q}_p \sigma^{\mu\nu} d_r) \tau^I \varphi W_{\mu\nu}^I$	$Q_{\varphi d}$	$(\varphi^\dagger i \overleftrightarrow{D}_\mu \varphi)(\bar{d}_p \gamma^\mu d_r)$
$Q_{\varphi \tilde{W}B}$	$\varphi^\dagger \tau^I \varphi \tilde{W}_{\mu\nu}^I B^{\mu\nu}$	Q_{dB}	$(\bar{q}_p \sigma^{\mu\nu} d_r) \varphi B_{\mu\nu}$	$Q_{\varphi ud}$	$i(\tilde{\varphi}^\dagger D_\mu \varphi)(\bar{u}_p \gamma^\mu d_r)$

$$(\phi^\dagger D_\mu \phi)(\bar{\psi} \gamma^\mu \psi)$$





Current operators

[Grzadkowski et al.; JHEP 10 (2010) 085]

X^3		φ^6 and $\varphi^4 D^2$		$\psi^2 \varphi^3$	
Q_G	$f^{ABC} G_\mu^{A\nu} G_\nu^{B\rho} G_\rho^{C\mu}$	Q_φ	$(\varphi^\dagger \varphi)^3$	$Q_{e\varphi}$	$(\varphi^\dagger \varphi)(\bar{l}_p e_r \varphi)$
$Q_{\tilde{G}}$	$f^{ABC} \tilde{G}_\mu^{A\nu} G_\nu^{B\rho} G_\rho^{C\mu}$	$Q_{\varphi\Box}$	$(\varphi^\dagger \varphi)\Box(\varphi^\dagger \varphi)$	$Q_{u\varphi}$	$(\varphi^\dagger \varphi)(\bar{q}_p u_r \tilde{\varphi})$
Q_W	$\varepsilon^{IJK} W_\mu^{I\nu} W_\nu^{J\rho} W_\rho^{K\mu}$	$Q_{\varphi D}$	$(\varphi^\dagger D^\mu \varphi)^* (\varphi^\dagger D_\mu \varphi)$	$Q_{d\varphi}$	$(\varphi^\dagger \varphi)(\bar{q}_p d_r \varphi)$
$Q_{\tilde{W}}$	$\varepsilon^{IJK} \tilde{W}_\mu^{I\nu} W_\nu^{J\rho} W_\rho^{K\mu}$				
$X^2 \varphi^2$		$\psi^2 X \varphi$		$\psi^2 \varphi^2 D$	
$Q_{\varphi G}$	$\varphi^\dagger \varphi G_{\mu\nu}^A G^{A\mu\nu}$	Q_{eW}	$(\bar{l}_p \sigma^{\mu\nu} e_r) \tau^I \varphi W_{\mu\nu}^I$	$Q_{\varphi l}^{(1)}$	$(\varphi^\dagger i \overleftrightarrow{D}_\mu \varphi)(\bar{l}_p \gamma^\mu l_r)$
$Q_{\varphi \tilde{G}}$	$\varphi^\dagger \varphi \tilde{G}_{\mu\nu}^A G^{A\mu\nu}$	Q_{eB}	$(\bar{l}_p \sigma^{\mu\nu} e_r) \varphi B_{\mu\nu}$	$Q_{\varphi l}^{(3)}$	$(\varphi^\dagger i \overleftrightarrow{D}_\mu^I \varphi)(\bar{l}_p \tau^I \gamma^\mu l_r)$
$Q_{\varphi W}$	$\varphi^\dagger \varphi W_{\mu\nu}^I W^{I\mu\nu}$	Q_{uG}	$(\bar{q}_p \sigma^{\mu\nu} T^A u_r) \tilde{\varphi} G_{\mu\nu}^A$	$Q_{\varphi e}$	$(\varphi^\dagger i \overleftrightarrow{D}_\mu \varphi)(\bar{e}_p \gamma^\mu e_r)$
$Q_{\varphi \tilde{W}}$	$\varphi^\dagger \varphi \tilde{W}_{\mu\nu}^I W^{I\mu\nu}$	Q_{uW}	$(\bar{q}_p \sigma^{\mu\nu} u_r) \tau^I \tilde{\varphi} W_{\mu\nu}^I$	$Q_{\varphi q}^{(1)}$	$(\varphi^\dagger i \overleftrightarrow{D}_\mu \varphi)(\bar{q}_p \gamma^\mu q_r)$
$Q_{\varphi B}$	$\varphi^\dagger \varphi B_{\mu\nu} B^{\mu\nu}$	Q_{uB}	$(\bar{q}_p \sigma^{\mu\nu} u_r) \tilde{\varphi} B_{\mu\nu}$	$Q_{\varphi q}^{(3)}$	$(\varphi^\dagger i \overleftrightarrow{D}_\mu^I \varphi)(\bar{q}_p \tau^I \gamma^\mu q_r)$
$Q_{\varphi \tilde{B}}$	$\varphi^\dagger \varphi \tilde{B}_{\mu\nu} B^{\mu\nu}$	Q_{dG}	$(\bar{q}_p \sigma^{\mu\nu} T^A d_r) \varphi G_{\mu\nu}^A$	$Q_{\varphi u}$	$(\varphi^\dagger i \overleftrightarrow{D}_\mu \varphi)(\bar{u}_p \gamma^\mu u_r)$
$Q_{\varphi WB}$	$\varphi^\dagger \tau^I \varphi W_{\mu\nu}^I B^{\mu\nu}$	Q_{dW}	$(\bar{q}_p \sigma^{\mu\nu} d_r) \tau^I \varphi W_{\mu\nu}^I$	$Q_{\varphi d}$	$(\varphi^\dagger i \overleftrightarrow{D}_\mu \varphi)(\bar{d}_p \gamma^\mu d_r)$
$Q_{\varphi \tilde{W}B}$	$\varphi^\dagger \tau^I \varphi \tilde{W}_{\mu\nu}^I B^{\mu\nu}$	Q_{dB}	$(\bar{q}_p \sigma^{\mu\nu} d_r) \varphi B_{\mu\nu}$	$Q_{\varphi ud}$	$i(\tilde{\varphi}^\dagger D_\mu \varphi)(\bar{u}_p \gamma^\mu d_r)$

$$(\phi^\dagger D_\mu \phi)(\bar{\psi} \gamma^\mu \psi)$$





EW Higgs production

Current operators


[Grzadkowski et al.; JHEP 10 (2010) 085]

X^3		φ^6 and $\varphi^4 D^2$		$\psi^2 \varphi^3$	
Q_G	$f^{ABC} G_\mu^{A\nu} G_\nu^{B\rho} G_\rho^{C\mu}$	Q_φ	$(\varphi^\dagger \varphi)^3$	$Q_{e\varphi}$	$(\varphi^\dagger \varphi)(\bar{l}_p e_r \varphi)$
$Q_{\tilde{G}}$	$f^{ABC} \tilde{G}_\mu^{A\nu} G_\nu^{B\rho} G_\rho^{C\mu}$	$Q_{\varphi\Box}$	$(\varphi^\dagger \varphi)\Box(\varphi^\dagger \varphi)$	$Q_{u\varphi}$	$(\varphi^\dagger \varphi)(\bar{q}_p u_r \tilde{\varphi})$
Q_W	$\varepsilon^{IJK} W_\mu^{I\nu} W_\nu^{J\rho} W_\rho^{K\mu}$	$Q_{\varphi D}$	$(\varphi^\dagger D^\mu \varphi)^* (\varphi^\dagger D_\mu \varphi)$	$Q_{d\varphi}$	$(\varphi^\dagger \varphi)(\bar{q}_p d_r \varphi)$
$Q_{\tilde{W}}$	$\varepsilon^{IJK} \tilde{W}_\mu^{I\nu} W_\nu^{J\rho} W_\rho^{K\mu}$				
$X^2 \varphi^2$		$\psi^2 X \varphi$		$\psi^2 \varphi^2 D$	
$Q_{\varphi G}$	$\varphi^\dagger \varphi G_{\mu\nu}^A G^{A\mu\nu}$	Q_{eW}	$(\bar{l}_p \sigma^{\mu\nu} e_r) \tau^I \varphi W_{\mu\nu}^I$	$Q_{\varphi l}^{(1)}$	$(\varphi^\dagger i \overleftrightarrow{D}_\mu \varphi)(\bar{l}_p \gamma^\mu l_r)$
$Q_{\varphi \tilde{G}}$	$\varphi^\dagger \varphi \tilde{G}_{\mu\nu}^A G^{A\mu\nu}$	Q_{eB}	$(\bar{l}_p \sigma^{\mu\nu} e_r) \varphi B_{\mu\nu}$	$Q_{\varphi l}^{(3)}$	$(\varphi^\dagger i \overleftrightarrow{D}_\mu^I \varphi)(\bar{l}_p \tau^I \gamma^\mu l_r)$
$Q_{\varphi W}$	$\varphi^\dagger \varphi W_{\mu\nu}^I W^{I\mu\nu}$	Q_{uG}	$(\bar{q}_p \sigma^{\mu\nu} T^A u_r) \tilde{\varphi} G_{\mu\nu}^A$	$Q_{\varphi e}$	$(\varphi^\dagger i \overleftrightarrow{D}_\mu \varphi)(\bar{e}_p \gamma^\mu e_r)$
$Q_{\varphi \tilde{W}}$	$\varphi^\dagger \varphi \tilde{W}_{\mu\nu}^I W^{I\mu\nu}$	Q_{uW}	$(\bar{q}_p \sigma^{\mu\nu} u_r) \tau^I \tilde{\varphi} W_{\mu\nu}^I$	$Q_{\varphi q}^{(1)}$	$(\varphi^\dagger i \overleftrightarrow{D}_\mu \varphi)(\bar{q}_p \gamma^\mu q_r)$
$Q_{\varphi B}$	$\varphi^\dagger \varphi B_{\mu\nu} B^{\mu\nu}$	Q_{uB}	$(\bar{q}_p \sigma^{\mu\nu} u_r) \tilde{\varphi} B_{\mu\nu}$	$Q_{\varphi q}^{(3)}$	$(\varphi^\dagger i \overleftrightarrow{D}_\mu^I \varphi)(\bar{q}_p \tau^I \gamma^\mu q_r)$
$Q_{\varphi \tilde{B}}$	$\varphi^\dagger \varphi \tilde{B}_{\mu\nu} B^{\mu\nu}$	Q_{dG}	$(\bar{q}_p \sigma^{\mu\nu} T^A d_r) \varphi G_{\mu\nu}^A$	$Q_{\varphi u}$	$(\varphi^\dagger i \overleftrightarrow{D}_\mu \varphi)(\bar{u}_p \gamma^\mu u_r)$
$Q_{\varphi WB}$	$\varphi^\dagger \tau^I \varphi W_{\mu\nu}^I B^{\mu\nu}$	Q_{dW}	$(\bar{q}_p \sigma^{\mu\nu} d_r) \tau^I \varphi W_{\mu\nu}^I$	$Q_{\varphi d}$	$(\varphi^\dagger i \overleftrightarrow{D}_\mu \varphi)(\bar{d}_p \gamma^\mu d_r)$
$Q_{\varphi \tilde{W}B}$	$\varphi^\dagger \tau^I \varphi \tilde{W}_{\mu\nu}^I B^{\mu\nu}$	Q_{dB}	$(\bar{q}_p \sigma^{\mu\nu} d_r) \varphi B_{\mu\nu}$	$Q_{\varphi ud}$	$i(\tilde{\varphi}^\dagger D_\mu \varphi)(\bar{u}_p \gamma^\mu d_r)$

$$(\phi^\dagger D_\mu \phi)(\bar{\psi} \gamma^\mu \psi)$$



EW Higgs production





Diboson production

Current operators


[Grzadkowski et al.; JHEP 10 (2010) 085]

X^3		φ^6 and $\varphi^4 D^2$		$\psi^2 \varphi^3$	
Q_G	$f^{ABC} G_{\mu}^{A\nu} G_{\nu}^{B\rho} G_{\rho}^{C\mu}$	Q_{φ}	$(\varphi^\dagger \varphi)^3$	$Q_{e\varphi}$	$(\varphi^\dagger \varphi)(\bar{l}_p e_r \varphi)$
$Q_{\tilde{G}}$	$f^{ABC} \tilde{G}_{\mu}^{A\nu} G_{\nu}^{B\rho} G_{\rho}^{C\mu}$	$Q_{\varphi\Box}$	$(\varphi^\dagger \varphi)\Box(\varphi^\dagger \varphi)$	$Q_{u\varphi}$	$(\varphi^\dagger \varphi)(\bar{q}_p u_r \tilde{\varphi})$
Q_W	$\varepsilon^{IJK} W_{\mu}^{I\nu} W_{\nu}^{J\rho} W_{\rho}^{K\mu}$	$Q_{\varphi D}$	$(\varphi^\dagger D^\mu \varphi)^* (\varphi^\dagger D_\mu \varphi)$	$Q_{d\varphi}$	$(\varphi^\dagger \varphi)(\bar{q}_p d_r \varphi)$
$Q_{\tilde{W}}$	$\varepsilon^{IJK} \tilde{W}_{\mu}^{I\nu} W_{\nu}^{J\rho} W_{\rho}^{K\mu}$				
$X^2 \varphi^2$		$\psi^2 X \varphi$		$\psi^2 \varphi^2 D$	
$Q_{\varphi G}$	$\varphi^\dagger \varphi G_{\mu\nu}^A G^{A\mu\nu}$	Q_{eW}	$(\bar{l}_p \sigma^{\mu\nu} e_r) \tau^I \varphi W_{\mu\nu}^I$	$Q_{\varphi l}^{(1)}$	$(\varphi^\dagger i \overleftrightarrow{D}_\mu \varphi)(\bar{l}_p \gamma^\mu l_r)$
$Q_{\varphi \tilde{G}}$	$\varphi^\dagger \varphi \tilde{G}_{\mu\nu}^A G^{A\mu\nu}$	Q_{eB}	$(\bar{l}_p \sigma^{\mu\nu} e_r) \varphi B_{\mu\nu}$	$Q_{\varphi l}^{(3)}$	$(\varphi^\dagger i \overleftrightarrow{D}_\mu^I \varphi)(\bar{l}_p \tau^I \gamma^\mu l_r)$
$Q_{\varphi W}$	$\varphi^\dagger \varphi W_{\mu\nu}^I W^{I\mu\nu}$	Q_{uG}	$(\bar{q}_p \sigma^{\mu\nu} T^A u_r) \tilde{\varphi} G_{\mu\nu}^A$	$Q_{\varphi e}$	$(\varphi^\dagger i \overleftrightarrow{D}_\mu \varphi)(\bar{e}_p \gamma^\mu e_r)$
$Q_{\varphi \tilde{W}}$	$\varphi^\dagger \varphi \tilde{W}_{\mu\nu}^I W^{I\mu\nu}$	Q_{uW}	$(\bar{q}_p \sigma^{\mu\nu} u_r) \tau^I \tilde{\varphi} W_{\mu\nu}^I$	$Q_{\varphi q}^{(1)}$	$(\varphi^\dagger i \overleftrightarrow{D}_\mu \varphi)(\bar{q}_p \gamma^\mu q_r)$
$Q_{\varphi B}$	$\varphi^\dagger \varphi B_{\mu\nu} B^{\mu\nu}$	Q_{uB}	$(\bar{q}_p \sigma^{\mu\nu} u_r) \tilde{\varphi} B_{\mu\nu}$	$Q_{\varphi q}^{(3)}$	$(\varphi^\dagger i \overleftrightarrow{D}_\mu^I \varphi)(\bar{q}_p \tau^I \gamma^\mu q_r)$
$Q_{\varphi \tilde{B}}$	$\varphi^\dagger \varphi \tilde{B}_{\mu\nu} B^{\mu\nu}$	Q_{dG}	$(\bar{q}_p \sigma^{\mu\nu} T^A d_r) \varphi G_{\mu\nu}^A$	$Q_{\varphi u}$	$(\varphi^\dagger i \overleftrightarrow{D}_\mu \varphi)(\bar{u}_p \gamma^\mu u_r)$
$Q_{\varphi WB}$	$\varphi^\dagger \tau^I \varphi W_{\mu\nu}^I B^{\mu\nu}$	Q_{dW}	$(\bar{q}_p \sigma^{\mu\nu} d_r) \tau^I \varphi W_{\mu\nu}^I$	$Q_{\varphi d}$	$(\varphi^\dagger i \overleftrightarrow{D}_\mu \varphi)(\bar{d}_p \gamma^\mu d_r)$
$Q_{\varphi \tilde{W}B}$	$\varphi^\dagger \tau^I \varphi \tilde{W}_{\mu\nu}^I B^{\mu\nu}$	Q_{dB}	$(\bar{q}_p \sigma^{\mu\nu} d_r) \varphi B_{\mu\nu}$	$Q_{\varphi ud}$	$i(\tilde{\varphi}^\dagger D_\mu \varphi)(\bar{u}_p \gamma^\mu d_r)$


$$(\phi^\dagger D_\mu \phi)(\bar{\psi} \gamma^\mu \psi)$$



EW Higgs production




Diboson production



Precision EW on the Z peak (LEP)

Interplay



[Ellis, Madigan, KM, Sanz & You; JHEP 04 (2021) 279]

Status in a nutshell

Global new physics searches via high precision/energy

- **Z & W-pole data:** handle on the EW gauge sector [Han & Skiba; PRD 71 (2005) 075009]
[Falkowski & Riva; JHEP 02 (2015) 039]
- **LHC:** thriving Higgs & top programmes
- Probing gauge interactions at high energy (**VV, VBS, VVV, ...**)

Status in a nutshell

Global new physics searches via high precision/energy

- **Z & W-pole data:** handle on the EW gauge sector [Han & Skiba; PRD 71 (2005) 075009]
[Falkowski & Riva; JHEP 02 (2015) 039]
- **LHC:** thriving Higgs & top programmes
- Probing gauge interactions at high energy (**VV, VBS, VVV, ...**)

How much cross-talk? Where does being global matter?

Status in a nutshell

Global new physics searches via high precision/energy

- **Z & W-pole data:** handle on the EW gauge sector [Han & Skiba; PRD 71 (2005) 075009]
[Falkowski & Riva; JHEP 02 (2015) 039]
- **LHC:** thriving Higgs & top programmes
- Probing gauge interactions at high energy (**VV, VBS, VVV, ...**)

How much cross-talk? Where does being global matter?

We know that Higgs data greatly complements LEP

- Access **unconstrained directions** in parameter space
- Allows for a **closed fit** to flavor-universal SMEFT
- Crucial to combine EWPO, Diboson & Higgs data

[Corbett et al.; PRD 87 (2013) 015022]


[Pomarol & Riva; JHEP 01 (2014) 151]

[Ellis, Sanz & You; JHEP 03 (2015) 157]

[Biekötter Corbett & Plehn; SciPost Phys 6 (2019) 6, 064]

[Anisha et al; arXiv:2111.05876].....

[Ellis et al.; JHEP 06
(2018) 146]




Top/Higgs interplay

$C_{\phi G}$ Point-like

$C_{t\phi}$ Yukawa

C_{tG} Dipole




Top/Higgs interplay

$C_{\phi G}$ Point-like

$C_{t\phi}$ Yukawa

C_{tG} Dipole



$gg \rightarrow h$ is well measured now...

⚠ Can't exclude top partners/anomalous Yukawa


- Degeneracy in coefficient/model space

Top/Higgs interplay

$C_{\phi G}$ Point-like

$C_{t\phi}$ Yukawa

C_{tG} Dipole



$gg \rightarrow h$ is well measured now...

⚠ Can't exclude top partners/anomalous Yukawa

- Degeneracy in coefficient/model space


Need more data!

Top/Higgs interplay

$C_{\phi G}$ Point-like

$C_{t\phi}$ Yukawa

C_{tG} Dipole



$gg \rightarrow h$ is well measured now...



⚠ Can't exclude top partners/anomalous Yukawa

- Degeneracy in coefficient/model space

Need more data!

- $t\bar{t}H$ production for direct Yukawa measurement
- $t\bar{t}$ data to constrain dipole

[Maltoni, Vryonidou &




Top/Higgs interplay

[Maltoni, Vryonidou &

$C_{\phi G}$ Point-like

$C_{t\phi}$ Yukawa

C_{tG} Dipole



$gg \rightarrow h$ is well measured now...

⚠ Can't exclude top partners/anomalous Yukawa




- Degeneracy in coefficient/model space

Need more data!

- $t\bar{t}H$ production for direct Yukawa measurement
- $t\bar{t}$ data to constrain dipole

Several other new interactions can affect $t\bar{t}$

- Notably $q\bar{q}t\bar{t}$ operators, of which there are many (14)
- To what extent do these limit ultimate NP sensitivity in top/Higgs sector?



Answer: do the fit

Top, Higgs, Diboson and Electroweak Fit to the
Standard Model Effective Field Theory

John Ellis,^{a,b,c} Maeve Madigan,^d Ken Mimasu,^a Veronica Sanz^{e,f} and Tevong You^{b,d,g}

[JHEP 04 (2021) 279]

Answer: do the fit

Top, Higgs, Diboson and Electroweak Fit to the Standard Model Effective Field Theory

John Ellis,^{a,b,c} Maeve Madigan,^d Ken Mimasu,^a Veronica Sanz^{e,f} and Tevong You^{b,d,g} [*JHEP* 04 (2021) 279]

Global SMEFT interpretation of 4 categories of data

- 14 • Electroweak Precision Observables (EWPO): Z-pole & W-mass
- 118 • LEP2 & LHC diboson production: differential WW, WZ, Zjj
- 72 • Higgs measurements: signal strengths & STXS
- 137 • Top data: single-top, ttbar & asymmetries, ttV, tZ, tW

Based on

[Ellis et al.; JHEP 06 (2018) 146]

Big thanks to authors of SMEFiT analysis [JHEP 04 (2019) 100] for sharing some of their top predictions

Answer: do the fit

Top, Higgs, Diboson and Electroweak Fit to the Standard Model Effective Field Theory

John Ellis,^{a,b,c} Maeve Madigan,^d Ken Mimasu,^a Veronica Sanz^{e,f} and Tevong You^{b,d,g} [*JHEP* 04 (2021) 279]

Global SMEFT interpretation of 4 categories of data

- 14 • Electroweak Precision Observables (EWPO): Z-pole & W-mass
- 118 • LEP2 & LHC diboson production: differential WW, WZ, Zjj
- 72 • Higgs measurements: signal strengths & STXS
- 137 • Top data: single-top, ttbar & asymmetries, ttV, tZ, tW

Based on
[Ellis et al.; *JHEP* 06 (2018) 146]

Big thanks to authors of
SMEFiT analysis
[*JHEP* 04 (2019) 100]
for sharing some of their
top predictions

341 measurements across categories

- Chosen to be statistically independent & maximise reach
- Correlations included when publicly available (mostly are)

Answer: do the fit

Top, Higgs, Diboson and Electroweak Fit to the Standard Model Effective Field Theory

John Ellis,^{a,b,c} Maeve Madigan,^d Ken Mimasu,^a Veronica Sanz^{e,f} and Tevong You^{b,d,g} [*JHEP* 04 (2021) 279]

Global SMEFT interpretation of 4 categories of data

Based on

- 14 • **Electroweak Precision Observables (EWPO):** Z-pole & W-mass [*Ellis et al.; JHEP* 06 (2018) 146]
- 118 • **LEP2 & LHC diboson production:** differential WW, WZ, Zjj
- 72 • **Higgs measurements:** signal strengths & STXS
- 137 • **Top data:** single-top, ttbar & asymmetries, ttV, tZ, tW

*Big thanks to authors of SMEFiT analysis [*JHEP* 04 (2019) 100] for sharing some of their top predictions*

341 measurements across categories

- Chosen to be statistically independent & maximise reach
- Correlations included when publicly available (mostly are)





Linear EFT approximation:
$$\mu_X \equiv \frac{X}{X_{SM}} = 1 + \sum_i a_i^X \frac{C_i}{\Lambda^2} + \mathcal{O}\left(\frac{1}{\Lambda^4}\right)$$

Degrees of freedom

EWPO:	$\mathcal{O}_{HWB}, \mathcal{O}_{HD}, \mathcal{O}_{ll}, \mathcal{O}_{Hl}^{(3)}, \mathcal{O}_{Hl}^{(1)}, \mathcal{O}_{He}, \mathcal{O}_{Hq}^{(3)}, \mathcal{O}_{Hq}^{(1)}, \mathcal{O}_{Hd}, \mathcal{O}_{Hu}$	
Bosonic:	$\mathcal{O}_{H\Box}, \mathcal{O}_{HG}, \mathcal{O}_{HW}, \mathcal{O}_{HB}, \mathcal{O}_W, \mathcal{O}_G,$	
Yukawa:	$\mathcal{O}_{\tau H}, \mathcal{O}_{\mu H}, \mathcal{O}_{bH}, \mathcal{O}_{tH},$	20
Top 2F:	$\mathcal{O}_{HQ}^{(3)}, \mathcal{O}_{HQ}^{(1)}, \mathcal{O}_{Ht}, \mathcal{O}_{tG}, \mathcal{O}_{tW}, \mathcal{O}_{tB},$	
Top 4F:	$\mathcal{O}_{Qq}^{3,1}, \mathcal{O}_{Qq}^{3,8}, \mathcal{O}_{Qq}^{1,8}, \mathcal{O}_{Qu}^8, \mathcal{O}_{Qd}^8, \mathcal{O}_{tQ}^8, \mathcal{O}_{tu}^8, \mathcal{O}_{td}^8.$	+14

Degrees of freedom



EWPO:	$\mathcal{O}_{HWB}, \mathcal{O}_{HD}, \mathcal{O}_{ll}, \mathcal{O}_{Hl}^{(3)}, \mathcal{O}_{Hl}^{(1)}, \mathcal{O}_{He}, \mathcal{O}_{Hq}^{(3)}, \mathcal{O}_{Hq}^{(1)}, \mathcal{O}_{Hd}, \mathcal{O}_{Hu}$	
Bosonic:	$\mathcal{O}_{H\Box}, \mathcal{O}_{HG}, \mathcal{O}_{HW}, \mathcal{O}_{HB}, \mathcal{O}_W, \mathcal{O}_G$	
Yukawa:	$\mathcal{O}_{\tau H}, \mathcal{O}_{\mu H}, \mathcal{O}_{bH}, \mathcal{O}_{tH}$	20
Top 2F:	$\mathcal{O}_{HQ}^{(3)}, \mathcal{O}_{HQ}^{(1)}, \mathcal{O}_{Ht}, \mathcal{O}_{tG}, \mathcal{O}_{tW}, \mathcal{O}_{tB}$	
Top 4F:	$\mathcal{O}_{Qq}^{3,1}, \mathcal{O}_{Qq}^{3,8}, \mathcal{O}_{Qq}^{1,8}, \mathcal{O}_{Qu}^8, \mathcal{O}_{Qd}^8, \mathcal{O}_{tQ}^8, \mathcal{O}_{tu}^8, \mathcal{O}_{td}^8$	+14



Individual limits: $U(3)^5$

2018 data: [Ellis et al.; JHEP 06 (2018) 146]



$SU(3)^5$: EWPO + Diboson + Higgs



Individual limits: $U(3)^5$

2018 data: [Ellis et al.; JHEP 06 (2018) 146]



$SU(3)^5$: EWPO + Diboson + Higgs



Individual limits: $U(3)^5$

2018 data: [Ellis et al.; JHEP 06 (2018) 146]



$SU(3)^5$: EWPO + Diboson + Higgs



Individual limits: $U(3)^5$



2018 data: [Ellis et al.; JHEP 06 (2018) 146]

$SU(3)^5$: EWPO + Diboson + Higgs





Marginalised limits: $U(3)^5$

2018 \Rightarrow 2020: only LHC data changed





Marginalised limits: $U(3)^5$

2018 \Rightarrow 2020: only LHC data changed





Marginalised limits: $U(3)^5$

2018 \Rightarrow 2020: only LHC data changed




Marginalised limits: $U(3)^5$

2018 \Rightarrow 2020: only LHC data changed



Top-Higgs interplay








2D constraints



Top-Higgs interplay

2D constraints

- $ggF/t\bar{t}H$ complementarity for (C_{HG}, C_{tH})
- H+jets STXS & $t\bar{t}V$ not yet competitive
- Strong impact of $t\bar{t}$ evident for (C_{tG}, C_G)
- Tension with SM $\sim 2\sigma$
- Significant correlations remain
- Large marginalisation effects



Point-like



Yukawa










Dipole

Top-Higgs interplay

2D constraints

- $ggF/t\bar{t}H$ complementarity for (C_{HG}, C_{tH})
- H+jets STXS & $t\bar{t}V$ not yet competitive
- Strong impact of $t\bar{t}$ evident for (C_{tG}, C_G)
- Tension with SM $\sim 2\sigma$
- Significant correlations remain
- Large marginalisation effects



What is the concrete impact of 4F?

4F impact

Fit to 'Higgs-only' subspace







$$C_{H\Box}, C_{HG}, C_{HW}, C_{HB}, C_{tH}, C_{bH}, C_{\tau H}, C_{\mu H} \\ + C_{tG} \text{ \& } C_G$$

- Allow a closed fit to Higgs data only
- Emphasises impact of $t\bar{t}H$ & $t\bar{t}$


Marginalised

Marginalised 95% C. L.


- Higgs data (no $t\bar{t}H$)
- Higgs data
- Higgs & Top data
- Higgs & Top data (+4F)
- + SM



Point-like



Yukawa



Dipole

4F impact

Fit to 'Higgs-only' subspace

$$C_{H\Box}, C_{HG}, C_{HW}, C_{HB}, C_{tH}, C_{bH}, C_{\tau H}, C_{\mu H} \\ + C_{tG} \text{ \& } C_G$$

- Allow a closed fit to Higgs data only
- Emphasises impact of $t\bar{t}H$ & $t\bar{t}$

Now add in $t\bar{t}$ 4F operators

$$+ C_{Qq}^{3,8}, C_{Qq}^{1,8}, C_{Qu}^8, C_{Qd}^8, C_{tq}^8, C_{tu}^8, C_{td}^8$$

- Relatively mild impact
- Preferred $t\bar{t}$ phase space is different

$$C_{tG} : \text{low } m_{t\bar{t}}$$

$$4F : \text{high } m_{t\bar{t}}$$







- Able to constrain them independently

[ATLAS-CONF-2021-031]

Marginalised

Marginalised 95% C. L.

- Higgs data (no $t\bar{t}H$)
- Higgs data
- Higgs & Top data
- Higgs & Top data (+4F)
- + SM



Point-like

Yukawa


Dipole

More top/Higgs interplay

[Ethier et al.; JHEP 11 (2021) 089]

Fisher Information:

*Hessian of Log-likelihood
at the best-fit point*




More top/Higgs interplay

[Ethier et al.; JHEP 11 (2021) 089]

Fisher Information:

*Hessian of Log-likelihood
at the best-fit point*




More top/Higgs interplay


[Ethier et al.; JHEP 11 (2021) 089]

Fisher Information:

*Hessian of Log-likelihood
at the best-fit point*



4F operators:
mostly top data




More top/Higgs interplay

[Ethier et al.; JHEP 11 (2021) 089]



Fisher Information:

Hessian of Log-likelihood
at the best-fit point



4F operators:
mostly top data

Normalized Fisher Value




Yukawa &
Chromo-dipole

More top/Higgs interplay

[Ethier et al.; JHEP 11 (2021) 089]




Fisher Information:

Hessian of Log-likelihood
at the best-fit point





4F operators:
mostly top data

Normalized Fisher Value




Full fit: marginalised



Correlations

Block diagonal:
correlations *within*
'sector'

Block off-diagonal:
correlations *among*
'sectors'




Correlations

Block diagonal:
correlations *within*
'sector'

Block off-diagonal:
correlations *among*
'sectors'

EWPO & top
~uncorrelated



Correlations


Block diagonal:
correlations *within*
'sector'

Block off-diagonal:
correlations *among*
'sectors'

EWPO & top
~uncorrelated

EWPO-Higgs
 $C_{HB}, C_{HW}, C_{H\Box}$
& Yukawa
with EWPO

Higgs precision
rivalling LEP



Correlations

Block diagonal:
correlations *within*
'sector'


Block off-diagonal:
correlations *among*
'sectors'

EWPO & top
~uncorrelated

EWPO-Higgs
 $C_{HB}, C_{HW}, C_{H\Box}$
& Yukawa
with EWPO


Higgs precision
rivalling LEP

Top-Higgs
 C_{HG}, C_G, C_{tH}
with 4F




PCA

2σ bound on Λ_i , $a_{ij}c_j = 1$





Relative constraining power (%)



PCA

2σ bound on $\Lambda_i, a_{ij}c_j = 1$



Relative constraining power (%)

6	-	15	29	50	-	-	-	-	-
35	-	13	28	23	-	-	-	-	-
59	-	7	16	18	-	-	-	-	-
-	-	5	94	1	-	-	-	-	-
99	-	-	-	-	-	-	-	-	-
95	-	-	-	3	-	-	-	-	-
99	-	-	-	-	-	-	-	-	-
-	-	21	45	33	-	-	-	-	-
74	-	2	4	14	6	-	-	-	-
18	-	8	19	55	-	-	-	-	-
73	22	-	-	-	3	-	-	-	-
7	2	5	53	4	29	-	-	-	-
-	-	-	3	-	-	-	-	96	-
7	-	2	39	2	46	-	-	3	-
-	-	-	10	4	-	-	57	-	28
94	2	-	-	-	2	-	-	-	-
-	2	-	-	-	11	86	-	-	-
65	14	2	4	10	4	1	-	-	-
-	-	-	-	-	-	-	99	-	-
-	-	-	-	-	-	-	84	15	-
-	-	-	13	6	-	-	48	-	32
90	5	-	-	1	2	-	-	-	-
-	-	-	5	2	-	-	79	-	13
-	-	-	-	-	-	-	3	78	18
2	3	27	40	12	6	-	4	-	4
-	-	1	2	-	-	-	15	4	77
4	4	14	33	23	7	1	12	-	2
-	-	1	3	1	-	-	4	-	89
-	-	3	10	4	-	-	28	2	53
1	-	8	27	16	2	-	32	-	12
-	-	-	-	-	-	-	58	-	41
-	-	-	-	-	-	-	90	-	10
-	-	-	-	-	-	-	61	-	38
-	-	-	-	-	-	-	95	-	5

Higgs

Run 1, Hf
Run 2, Hf
STXS

BSM implications

SMEFT-UV connection is model dependent by construction

BSM implications

SMEFT-UV connection is model dependent by construction

- New physics implications & EFT validity are ***a posteriori***
- Depends on **sensitivity** & **energy scale** probed by data
- Bottom-up philosophy: new physics scale unknown

BSM implications

SMEFT-UV connection is model dependent by construction

- New physics implications & EFT validity are ***a posteriori***
- Depends on **sensitivity** & **energy scale** probed by data
- Bottom-up philosophy: new physics scale unknown

arbitrary dimensionful parameter $\frac{c_S}{\Lambda^2} = \frac{\lambda^2}{M^2}$ *coupling/mass scale of new physics*


BSM implications

SMEFT-UV connection is model dependent by construction

- New physics implications & EFT validity are ***a posteriori***
- Depends on **sensitivity** & **energy scale** probed by data
- Bottom-up philosophy: new physics scale unknown

arbitrary dimensionful parameter $\frac{c_S}{\Lambda^2} = \frac{\lambda^2}{M^2}$ *coupling/mass scale of new physics*

constraint: $c/\Lambda^2 < X$




BSM implications

SMEFT-UV connection is model dependent by construction

- New physics implications & EFT validity are ***a posteriori***
- Depends on **sensitivity** & **energy scale** probed by data
- Bottom-up philosophy: new physics scale unknown

arbitrary dimensionful parameter $\frac{c_S}{\Lambda^2} = \frac{\lambda^2}{M^2}$ coupling/mass scale of new physics

constraint: $c/\Lambda^2 < X$



Difficult to address in a general way


BSM implications

SMEFT-UV connection is model dependent by construction

- New physics implications & EFT validity are ***a posteriori***
- Depends on **sensitivity** & **energy scale** probed by data
- Bottom-up philosophy: new physics scale unknown

arbitrary dimensionful parameter $\frac{c_S}{\Lambda^2} = \frac{\lambda^2}{M^2}$ coupling/mass scale of new physics

constraint: $c/\Lambda^2 < X$



Difficult to address in a general way

- Today we are probing TeV scale new physics
- Hierarchies in sensitivity EWPO \gtrsim Higgs $>$ top (EW)


BSM implications

SMEFT-UV connection is model dependent by construction

- New physics implications & EFT validity are ***a posteriori***
- Depends on **sensitivity** & **energy scale** probed by data
- Bottom-up philosophy: new physics scale unknown

arbitrary dimensionful parameter $\frac{c_S}{\Lambda^2} = \frac{\lambda^2}{M^2}$ coupling/mass scale of new physics


constraint: $c/\Lambda^2 < X$



Difficult to address in a general way

- Today we are probing TeV scale new physics
- Hierarchies in sensitivity EWPO \gtrsim Higgs $>$ top (EW)
- Moderate-to-strong coupling scenarios most safe
- **Generic NP in loops** looks challenging for the LHC

BSM implications



Individual/marginalised = optimistic/pessimistic

- Real models should lie somewhere in between
- Less underlying parameters - more correlations
- Need to 're-run' the fits to infer on underlying model parameters

Single field extensions

Name	Spin	SU(3)	SU(2)	U(1)	Param.	Name	Spin	SU(3)	SU(2)	U(1)	Param.	
S	0	1	1	0	(M_S, κ_S)	Δ_1	$\frac{1}{2}$	1	2	$-\frac{1}{2}$	$(M_{\Delta_1}, \lambda_{\Delta_1})$	VLL
S_1	0	1	1	1	(M_{S_1}, y_{S_1})	Δ_3	$\frac{1}{2}$	1	2	$-\frac{1}{2}$	$(M_{\Delta_3}, \lambda_{\Delta_3})$	
φ	0	1	2	$\frac{1}{2}$	$(M_\varphi, Z_6 \cos \beta)$	Σ	$\frac{1}{2}$	1	3	0	$(M_\Sigma, \lambda_\Sigma)$	
Ξ	0	1	3	0	(M_Ξ, κ_Ξ)	Σ_1	$\frac{1}{2}$	1	3	-1	$(M_{\Sigma_1}, \lambda_{\Sigma_1})$	
Ξ_1	0	1	3	1	$(M_{\Xi_1}, \kappa_{\Xi_1})$	U	$\frac{1}{2}$	3	1	$\frac{2}{3}$	(M_U, λ_U)	VLL
B	1	1	1	0	(M_B, \hat{g}_H^B)	D	$\frac{1}{2}$	3	1	$-\frac{1}{3}$	(M_D, λ_D)	
B_1	1	1	1	1	(M_{B_1}, g_{B_1})	Q_1	$\frac{1}{2}$	3	2	$\frac{1}{6}$	(M_{Q_1}, λ_{Q_1})	
W	1	1	3	0	(M_W, \hat{g}_H^W)	Q_5	$\frac{1}{2}$	3	2	$-\frac{5}{6}$	(M_{Q_5}, λ_{Q_5})	
W_1	1	1	3	1	$(M_{W_1}, \hat{g}_{W_1}^\varphi)$	Q_7	$\frac{1}{2}$	3	2	$\frac{7}{6}$	(M_{Q_7}, λ_{Q_7})	
N	$\frac{1}{2}$	1	1	0	(M_N, λ_N)	T_1	$\frac{1}{2}$	3	3	$-\frac{1}{3}$	(M_{T_1}, λ_{T_1})	VLL
E	$\frac{1}{2}$	1	1	-1	(M_E, λ_E)	T_2	$\frac{1}{2}$	3	3	$\frac{2}{3}$	(M_{T_2}, λ_{T_2})	
T	$\frac{1}{2}$	3	1	$\frac{2}{3}$	(M_T, s_L^t)	TB	$\frac{1}{2}$	3	2	$\frac{1}{6}$	$(M_{TB}, s_L^{t,b})$	VLQ

Single field extensions

Name	Spin	SU(3)	SU(2)	U(1)	Param.	Name	Spin	SU(3)	SU(2)	U(1)	Param.
S	0	1	1	0	(M_S, κ_S)	Δ_1	$\frac{1}{2}$	1	2	$-\frac{1}{2}$	$(M_{\Delta_1}, \lambda_{\Delta_1})$
S_1	0	1	1	1	(M_{S_1}, y_{S_1})	Δ_3	$\frac{1}{2}$	1	2	$-\frac{1}{2}$	$(M_{\Delta_3}, \lambda_{\Delta_3})$
φ	0	1	2	$\frac{1}{2}$	$(M_\varphi, Z_6 \cos \beta)$	Σ	$\frac{1}{2}$	1	3	0	$(M_\Sigma, \lambda_\Sigma)$
Ξ	0	1	3	0	(M_Ξ, κ_Ξ)	Σ_1	$\frac{1}{2}$	1	3	-1	$(M_{\Sigma_1}, \lambda_{\Sigma_1})$
Ξ_1	0	1	3	1	$(M_{\Xi_1}, \kappa_{\Xi_1})$	U	$\frac{1}{2}$	3	1	$\frac{2}{3}$	(M_U, λ_U)
B	1	1	1	0	(M_B, \hat{g}_H^B)	D	$\frac{1}{2}$	3	1	$-\frac{1}{3}$	(M_D, λ_D)
B_1	1	1	1	1	(M_{B_1}, g_{B_1})	Q_1	$\frac{1}{2}$	3	2	$\frac{1}{6}$	(M_{Q_1}, λ_{Q_1})
W	1	1	3	0	(M_W, \hat{g}_H^W)	Q_5	$\frac{1}{2}$	3	2	$-\frac{5}{6}$	(M_{Q_5}, λ_{Q_5})
W_1	1	1	3	1	$(M_{W_1}, \hat{g}_{W_1}^\varphi)$	Q_7	$\frac{1}{2}$	3	2	$\frac{7}{6}$	(M_{Q_7}, λ_{Q_7})
N	$\frac{1}{2}$	1	1	0	(M_N, λ_N)	T_1	$\frac{1}{2}$	3	3	$-\frac{1}{3}$	(M_{T_1}, λ_{T_1})
E	$\frac{1}{2}$	1	1	-1	(M_E, λ_E)	T_2	$\frac{1}{2}$	3	3	$\frac{2}{3}$	(M_{T_2}, λ_{T_2})
T	$\frac{1}{2}$	3	1	$\frac{2}{3}$	(M_T, s_L^t)	TB	$\frac{1}{2}$	3	2	$\frac{1}{6}$	$(M_{TB}, s_L^{t,b})$

Scalars

VLL

Z'

W'

VLQ

VLL

VLQ

Considered single field extensions of the SM

Single field extensions

Name	Spin	SU(3)	SU(2)	U(1)	Param.	Name	Spin	SU(3)	SU(2)	U(1)	Param.	
S	0	1	1	0	(M_S, κ_S)	Δ_1	$\frac{1}{2}$	1	2	$-\frac{1}{2}$	$(M_{\Delta_1}, \lambda_{\Delta_1})$	VLL
S_1	0	1	1	1	(M_{S_1}, y_{S_1})	Δ_3	$\frac{1}{2}$	1	2	$-\frac{1}{2}$	$(M_{\Delta_3}, \lambda_{\Delta_3})$	
φ	0	1	2	$\frac{1}{2}$	$(M_\varphi, Z_6 \cos \beta)$	Σ	$\frac{1}{2}$	1	3	0	$(M_\Sigma, \lambda_\Sigma)$	
Ξ	0	1	3	0	(M_Ξ, κ_Ξ)	Σ_1	$\frac{1}{2}$	1	3	-1	$(M_{\Sigma_1}, \lambda_{\Sigma_1})$	
Ξ_1	0	1	3	1	$(M_{\Xi_1}, \kappa_{\Xi_1})$	U	$\frac{1}{2}$	3	1	$\frac{2}{3}$	(M_U, λ_U)	VLL
B	1	1	1	0	(M_B, \hat{g}_H^B)	D	$\frac{1}{2}$	3	1	$-\frac{1}{3}$	(M_D, λ_D)	
B_1	1	1	1	1	(M_{B_1}, g_{B_1})	Q_1	$\frac{1}{2}$	3	2	$\frac{1}{6}$	(M_{Q_1}, λ_{Q_1})	
W	1	1	3	0	(M_W, \hat{g}_H^W)	Q_5	$\frac{1}{2}$	3	2	$-\frac{5}{6}$	(M_{Q_5}, λ_{Q_5})	
W_1	1	1	3	1	$(M_{W_1}, \hat{g}_{W_1}^\varphi)$	Q_7	$\frac{1}{2}$	3	2	$\frac{7}{6}$	(M_{Q_7}, λ_{Q_7})	VLQ
N	$\frac{1}{2}$	1	1	0	(M_N, λ_N)	T_1	$\frac{1}{2}$	3	3	$-\frac{1}{3}$	(M_{T_1}, λ_{T_1})	
E	$\frac{1}{2}$	1	1	-1	(M_E, λ_E)	T_2	$\frac{1}{2}$	3	3	$\frac{2}{3}$	(M_{T_2}, λ_{T_2})	VLQ
T	$\frac{1}{2}$	3	1	$\frac{2}{3}$	(M_T, s_L^t)	TB	$\frac{1}{2}$	3	2	$\frac{1}{6}$	$(M_{TB}, s_L^{t,b})$	

Considered single field extensions of the SM

- Complete tree-level matching dictionary is known *[de Blas et al.; JHEP 03 (2018) 109]*

Single field extensions

Name	Spin	SU(3)	SU(2)	U(1)	Param.	Name	Spin	SU(3)	SU(2)	U(1)	Param.	
S	0	1	1	0	(M_S, κ_S)	Δ_1	$\frac{1}{2}$	1	2	$-\frac{1}{2}$	$(M_{\Delta_1}, \lambda_{\Delta_1})$	VLL
S_1	0	1	1	1	(M_{S_1}, y_{S_1})	Δ_3	$\frac{1}{2}$	1	2	$-\frac{1}{2}$	$(M_{\Delta_3}, \lambda_{\Delta_3})$	
φ	0	1	2	$\frac{1}{2}$	$(M_\varphi, Z_6 \cos \beta)$	Σ	$\frac{1}{2}$	1	3	0	$(M_\Sigma, \lambda_\Sigma)$	
Ξ	0	1	3	0	(M_Ξ, κ_Ξ)	Σ_1	$\frac{1}{2}$	1	3	-1	$(M_{\Sigma_1}, \lambda_{\Sigma_1})$	
Ξ_1	0	1	3	1	$(M_{\Xi_1}, \kappa_{\Xi_1})$	U	$\frac{1}{2}$	3	1	$\frac{2}{3}$	(M_U, λ_U)	VLQ
B	1	1	1	0	(M_B, \hat{g}_H^B)	D	$\frac{1}{2}$	3	1	$-\frac{1}{3}$	(M_D, λ_D)	
B_1	1	1	1	1	(M_{B_1}, g_{B_1})	Q_1	$\frac{1}{2}$	3	2	$\frac{1}{6}$	(M_{Q_1}, λ_{Q_1})	
W	1	1	3	0	(M_W, \hat{g}_H^W)	Q_5	$\frac{1}{2}$	3	2	$-\frac{5}{6}$	(M_{Q_5}, λ_{Q_5})	
W_1	1	1	3	1	$(M_{W_1}, \hat{g}_{W_1}^\varphi)$	Q_7	$\frac{1}{2}$	3	2	$\frac{7}{6}$	(M_{Q_7}, λ_{Q_7})	
N	$\frac{1}{2}$	1	1	0	(M_N, λ_N)	T_1	$\frac{1}{2}$	3	3	$-\frac{1}{3}$	(M_{T_1}, λ_{T_1})	VLL
E	$\frac{1}{2}$	1	1	-1	(M_E, λ_E)	T_2	$\frac{1}{2}$	3	3	$\frac{2}{3}$	(M_{T_2}, λ_{T_2})	
T	$\frac{1}{2}$	3	1	$\frac{2}{3}$	(M_T, s_L^t)	TB	$\frac{1}{2}$	3	2	$\frac{1}{6}$	$(M_{TB}, s_L^{t,b})$	VLQ

Considered single field extensions of the SM

- Complete tree-level matching dictionary is known *[de Blas et al.; JHEP 03 (2018) 109]*
- Interpret in terms of simplified 1 & 2 parameter versions of the models


One parameter models

Model	C_{HD}	C_{ll}	C_{Hl}^3	C_{Hl}^1	C_{He}	$C_{H\Box}$	$C_{\tau H}$	C_{tH}	C_{bH}
S						$-\frac{1}{2}$			
S_1		1							
Σ			$\frac{1}{16}$	$\frac{3}{16}$			$\frac{y_\tau}{4}$		
Σ_1			$-\frac{1}{16}$	$-\frac{3}{16}$			$\frac{y_\tau}{8}$		
N			$-\frac{1}{4}$	$\frac{1}{4}$					
E			$-\frac{1}{4}$	$-\frac{1}{4}$			$\frac{y_\tau}{2}$		
Δ_1					$\frac{1}{2}$		$\frac{y_\tau}{2}$		
Δ_3					$-\frac{1}{2}$		$\frac{y_\tau}{2}$		
B_1	1					$-\frac{1}{2}$	$-\frac{y_\tau}{2}$	$-\frac{y_t}{2}$	$-\frac{y_b}{2}$
Ξ	-2					$\frac{1}{2}$	y_τ	y_t	y_b
W_1	$-\frac{1}{4}$					$-\frac{1}{8}$	$-\frac{y_\tau}{8}$	$-\frac{y_t}{8}$	$-\frac{y_b}{8}$
φ							$-y_\tau$	$-y_t$	$-y_b$
$\{B, B_1\}$						$-\frac{3}{2}$	$-y_\tau$	$-y_t$	$-y_b$
$\{Q_1, Q_7\}$								y_t	

Model	C_{Hq}^3	C_{Hq}^1	$(C_{Hq}^3)_{33}$	$(C_{Hq}^1)_{33}$	C_{Hu}	C_{Hd}	C_{tH}	C_{bH}
U	$-\frac{1}{4}$	$\frac{1}{4}$	$-\frac{1}{4}$	$\frac{1}{4}$			$\frac{y_t}{2}$	
D	$-\frac{1}{4}$	$-\frac{1}{4}$	$-\frac{1}{4}$	$-\frac{1}{4}$				$\frac{y_b}{2}$
Q_5						$-\frac{1}{2}$		$\frac{y_b}{2}$
Q_7					$\frac{1}{2}$		$\frac{y_t}{2}$	
T_1	$-\frac{1}{16}$	$-\frac{3}{16}$	$-\frac{1}{16}$	$-\frac{3}{16}$			$\frac{y_t}{4}$	$\frac{y_b}{8}$
T_2	$-\frac{1}{16}$	$\frac{3}{16}$	$-\frac{1}{16}$	$\frac{3}{16}$			$\frac{y_t}{8}$	$\frac{y_b}{4}$
T			$-\frac{1}{2} \frac{M_T^2}{v^2}$	$\frac{1}{2} \frac{M_T^2}{v^2}$			$y_t \frac{M_T^2}{v^2}$	

$$\times \frac{\lambda^2}{M^2}$$


One parameter models



Pull-ology

Brute force: fit to all combinations of n-coefficients

$$P \equiv \sqrt{\chi_{SM}^2 - \chi_{fit}^2}$$




Pull-ology

Brute force: fit to all combinations of n-coefficients

$$P \equiv \sqrt{\chi_{SM}^2 - \chi_{fit}^2}$$

- Agnostic search for improved fit w.r.t SM
- NP hints could show up in this way
- Advantage of fast, linear fit method




Pull-ology

Brute force: fit to all combinations of n-coefficients

$$P \equiv \sqrt{\chi_{SM}^2 - \chi_{fit}^2}$$

- Agnostic search for improved fit w.r.t SM
- NP hints could show up in this way
- Advantage of fast, linear fit method
- Highlights tension in $t\bar{t}$ data
- No conclusive NP hints so far...



Conclusions

Presented first EWPO, Higgs, Diboson & Top fit in SMEFT

- Higgs data are driving global sensitivity beyond EWPO
- Top & Higgs sector are starting to talk to each other
- Interplay is robust against 4F operators in $t\bar{t}$

Analytic, linear analysis has many benefits

- Simple likelihood described by best fit+correlations, PCA exact
- Easy to interpret/combine with other likelihoods
- Fast: repeat for subsets, BSM interpretations

Drawbacks

- Potentially important quadratic effects, especially in top data
- Gaussian priors only, not really appropriate for th. errors

Outlook

Much more to be done

- Explore the likelihood further - how? opportunity for exp/th exchange
- Compare results to a quadratic fit to test validity
- SMEFT theory errors?
- Explore impact of new observables: VBS, VVV, rare top modes

Impact of loops

- Top operators in loops: Higgs decays + EWPO
- NLO corrections


BSM implications

- Go beyond 1 particle benchmarks towards realistic models
- Are there compelling top/Higgs scenarios that admit a valid EFT interpretation with LHC data?

Bonus: m_W interpretation



The result




$$\mathcal{O}_{HWB} \equiv H^\dagger \tau^I H W_{\mu\nu}^I B^{\mu\nu}, \quad \mathcal{O}_{HD} \equiv \left(H^\dagger D^\mu H \right)^* \left(H^\dagger D_\mu H \right)$$


$$\mathcal{O}_{\ell\ell} \equiv \left(\bar{\ell}_p \gamma_\mu \ell_r \right) \left(\bar{\ell}_s \gamma^\mu \ell_t \right), \quad \mathcal{O}_{H\ell}^{(3)} \equiv \left(H^\dagger i \overleftrightarrow{D}_\mu^I H \right) \left(\bar{\ell}_p \tau^I \gamma^\mu \ell_r \right)$$

$$\frac{\delta m_W^2}{m_W^2} = -\frac{\sin 2\theta_w}{\cos 2\theta_w} \frac{v^2}{4\Lambda^2} \left(\frac{\cos \theta_w}{\sin \theta_w} C_{HD} + \frac{\sin \theta_w}{\cos \theta_w} \left(4C_{H\ell}^{(3)} - 2C_{\ell\ell} \right) + 4C_{HWB} \right)$$


S & T parameters





Global fit (20 parameters)



Global fit (20 parameters)




Subsets



Single field extensions

Model	C_{HD}	C_{ll}	$C_{Hl}^{(3)}$	$C_{Hl}^{(1)}$	C_{He}	$C_{H\Box}$	$C_{\tau H}$	C_{tH}	C_{bH}
S_1		-1							
Σ			$\frac{1}{16}$	$\frac{3}{16}$			$\frac{y_\tau}{4}$		
Σ_1			$\frac{1}{16}$	$-\frac{3}{16}$			$\frac{y_\tau}{8}$		
N			$-\frac{1}{4}$	$\frac{1}{4}$					
E			$-\frac{1}{4}$	$-\frac{1}{4}$			$\frac{y_\tau}{2}$		
B_1	1					$-\frac{1}{2}$	$-\frac{y_\tau}{2}$	$-\frac{y_t}{2}$	$-\frac{y_b}{2}$
B	-2						$-y_\tau$	$-y_t$	$-y_b$
Ξ	$-2 \left(\frac{1}{M_\Xi}\right)^2$					$\frac{1}{2} \left(\frac{1}{M_\Xi}\right)^2$	$y_\tau \left(\frac{1}{M_\Xi}\right)^2$	$y_t \left(\frac{1}{M_\Xi}\right)^2$	$y_b \left(\frac{1}{M_\Xi}\right)^2$
W_1	$-\frac{1}{4}$					$-\frac{1}{8}$	$-\frac{y_\tau}{8}$	$-\frac{y_t}{8}$	$-\frac{y_b}{8}$
W	$\frac{1}{2}$					$-\frac{1}{2}$	$-y_\tau$	$-y_t$	$-y_b$



Backup



Data: EWPO & Diboson

EW precision observables	n_{obs}
Precision electroweak measurements on the Z resonance. $\Gamma_Z, \sigma_{\text{had.}}^0, R_\ell^0, A_{FB}^\ell, A_\ell(\text{SLD}), A_\ell(\text{Pt}), R_b^0, R_c^0, A_{FB}^b, A_{FB}^c, A_b$ & A_c	12
Combination of CDF and D0 W -Boson Mass Measurements	1
LHC run 1 W boson mass measurement by ATLAS	1
Diboson LEP & LHC	n_{obs}
$W^+ W^-$ angular distribution measurements at LEP II.	8
$W^+ W^-$ total cross section measurements at L3 in the $l\nu l\nu, l\nu qq$ & $qqqq$ final states for 8 energies	24
$W^+ W^-$ total cross section measurements at OPAL in the $l\nu l\nu, l\nu qq$ & $qqqq$ final states for 7 energies	21
$W^+ W^-$ total cross section measurements at ALEPH in the $l\nu l\nu, l\nu qq$ & $qqqq$ final states for 8 energies	21
ATLAS $W^+ W^-$ differential cross section in the $e\nu\mu\nu$ channel, $\frac{d\sigma}{dp_{\ell_1}^T}$, $p_T > 120$ GeV overflow bin	1
ATLAS $W^+ W^-$ fiducial differential cross section in the $e\nu\mu\nu$ channel, $\frac{d\sigma}{dp_{\ell_1}^T}$	14
ATLAS $W^\pm Z$ fiducial differential cross section in the $l^+ l^- l^\pm \nu$ channel, $\frac{d\sigma}{dp_Z^T}$	7
CMS $W^\pm Z$ normalised fiducial differential cross section in the $l^+ l^- l^\pm \nu$ channel, $\frac{1}{\sigma} \frac{d\sigma}{dp_Z^T}$	11
ATLAS Zjj fiducial differential cross section in the $l^+ l^-$ channel, $\frac{d\sigma}{d\Delta\varphi_{jj}}$	12

Data: Higgs

LHC Run 1 Higgs	n_{obs}
ATLAS and CMS LHC Run 1 combination of Higgs signal strengths. Production: ggF, VBF, ZH, WH & ttH Decay: $\gamma\gamma, ZZ, W^+W^-, \tau^+\tau^-$ & $b\bar{b}$	21
ATLAS inclusive $Z\gamma$ signal strength measurement	1
LHC Run 2 Higgs (new)	n_{obs}
ATLAS combination of signal strengths and stage 1.0 STXS in $H \rightarrow 4\ell$ including ratios of branching fractions to $\gamma\gamma, WW^*, \tau^+\tau^-$ & $b\bar{b}$ Signal strengths coarse STXS bins fine STXS bins	16 19 25
CMS LHC combination of Higgs signal strengths. Production: ggF, VBF, ZH, WH & ttH Decay: $\gamma\gamma, ZZ, W^+W^-, \tau^+\tau^-, b\bar{b}$ & $\mu^+\mu^-$	23
CMS stage 1.0 STXS measurements for $H \rightarrow \gamma\gamma$. 13 parameter fit 7 parameter fit	13 7
CMS stage 1.0 STXS measurements for $H \rightarrow \tau^+\tau^-$	9
CMS stage 1.1 STXS measurements for $H \rightarrow 4\ell$	19
CMS differential cross section measurements of inclusive Higgs production in the $WW^* \rightarrow \ell\nu\ell\nu$ final state. $\frac{d\sigma}{dn_{\text{jet}}}$ $\frac{d\sigma}{dp_H^T}$	5 6
ATLAS $H \rightarrow Z\gamma$ signal strength.	1
ATLAS $H \rightarrow \mu^+\mu^-$ signal strength.	1

Data: Tevatron, LHC Run 1 & 2 top

Tevatron & Run 1 top	n_{obs}	Ref.	Run 2 top	n_{obs}	Ref.
Tevatron combination of differential $t\bar{t}$ forward-backward asymmetry, $A_{FB}(m_{t\bar{t}})$.	4	[7]	CMS $t\bar{t}$ differential distributions in the dilepton channel. $\frac{d\sigma}{dm_{t\bar{t}}}$	6	[46, 50]
ATLAS $t\bar{t}$ differential distributions in the dilepton channel. $\frac{d\sigma}{dm_{t\bar{t}}}$	6	[31]	CMS $t\bar{t}$ differential distributions in the ℓ +jets channel. $\frac{d\sigma}{dm_{t\bar{t}}}$	10	[53]
ATLAS $t\bar{t}$ differential distributions in the ℓ +jets channel. $\frac{d\sigma}{dm_{t\bar{t}}}$ $\frac{d\sigma}{d y_{t\bar{t}} }$ $\frac{d\sigma}{dp_t^T}$ $\frac{d\sigma}{d y_t }$	7 5 8 5	[24]	ATLAS measurement of differential $t\bar{t}$ charge asymmetry, $A_C(m_{t\bar{t}})$.	5	[55]
CMS $t\bar{t}$ differential distributions in the ℓ +jets channel. $\frac{d\sigma}{dm_{t\bar{t}}}$ $\frac{d\sigma}{dy_{t\bar{t}}}$ $\frac{d\sigma}{dp_t^T}$ $\frac{d\sigma}{dy_t}$	7 10 8 10	[25, 34]	ATLAS $t\bar{t}W$ & $t\bar{t}Z$ cross section measurements. $\sigma_{t\bar{t}W} \sigma_{t\bar{t}Z}$	2	[58]
CMS measurement of differential $t\bar{t}$ charge asymmetry, $A_C(m_{t\bar{t}})$ in the dilepton channel.	3	[33]	CMS $t\bar{t}W$ & $t\bar{t}Z$ cross section measurements. $\sigma_{t\bar{t}W} \sigma_{t\bar{t}Z}$	1 1	[48]
ATLAS inclusive measurement $t\bar{t}$ charge asymmetry, $A_C(m_{t\bar{t}})$ in the dilepton channel.	1	[32]	CMS $t\bar{t}Z$ differential distributions. $\frac{d\sigma}{dp_z^T}$ $\frac{d\sigma}{d\cos\theta^*}$	4 4	[60]
ATLAS & CMS combination of differential $t\bar{t}$ charge asymmetry, $A_C(m_{t\bar{t}})$, in the ℓ +jets channel.	6	[38]	ATLAS $t\bar{t}\gamma$ differential distribution.	11	[62]
CMS $t\bar{t}$ double differential distributions in the dilepton channel. $\frac{d\sigma}{dm_{t\bar{t}}dy_t}$ $\frac{d\sigma}{dm_{t\bar{t}}dy_{t\bar{t}}}$ $\frac{d\sigma}{dm_{t\bar{t}}dp_{t\bar{t}}^T}$ $\frac{d\sigma}{dy_t dp_{t\bar{t}}^T}$	16 16 16 16	[18, 35]	CMS measurement of differential cross sections and charge ratios for t -channel single-top quark production. $\frac{d\sigma}{dp_{t+\bar{t}}^T}$ $R_t(p_{t+\bar{t}}^T)$	5 5	[56]
ATLAS & CMS Run 1 combination of W -boson helicity fractions in top decay. f_0, f_L & f_R	3	[40]	CMS measurement of t -channel single-top and anti-top cross sections. $\sigma_t, \sigma_{\bar{t}}, \sigma_{t+\bar{t}}$ & R_t .	4	[42]
ATLAS measurement of W -boson helicity fractions in top decay. f_0, f_L & f_R	3	[30]	CMS measurement of the t -channel single-top and anti-top cross sections. $\sigma_t \sigma_{\bar{t}} \sigma_{t+\bar{t}} R_t$.	1 1 1 1	[45]
CMS measurement of W -boson helicity fractions in top decay. f_0, f_L & f_R	3	[29]	CMS t -channel single-top differential distributions. $\frac{d\sigma}{dp_{t+\bar{t}}^T}$ $\frac{d\sigma}{d y_{t+\bar{t}} }$	4 4	[44]
ATLAS $t\bar{t}W$ & $t\bar{t}Z$ cross section measurements. $\sigma_{t\bar{t}W} \sigma_{t\bar{t}Z}$	2	[23]	ATLAS tW cross section measurement.	1	[43]
CMS $t\bar{t}W$ & $t\bar{t}Z$ cross section measurements. $\sigma_{t\bar{t}W} \sigma_{t\bar{t}Z}$	2	[26]	CMS tZ cross section measurement.	1	[47]
ATLAS $t\bar{t}\gamma$ cross section measurement in the ℓ + jets channel.	1	[36]	CMS tW cross section measurement.	1	[52]
CMS $t\bar{t}\gamma$ cross section measurement in the ℓ + jets channel.	1	[37]	ATLAS tZ cross section measurement.	1	[49]
ATLAS t -channel single-top differential distributions. $\frac{d\sigma}{dp_t^T}$ $\frac{d\sigma}{dp_{\bar{t}}^T}$ $\frac{d\sigma}{d y_t }$ $\frac{d\sigma}{d y_{\bar{t}} }$	4 4 4 5	[39]	CMS tZ ($Z \rightarrow \ell^+\ell^-$) cross section measurement	1	[54]
CMS s -channel single-top cross section measurement.	1	[28]	ATLAS four-top search in the multi-lepton and same-sign dilepton channels.	1	[63]
CMS t -channel single-top differential distributions. $\frac{d\sigma}{dp_{t+\bar{t}}^T}$ $\frac{d\sigma}{d y_{t+\bar{t}} }$	6 6	[19]	ATLAS four-top search in the single-lepton and opposite-sign dilepton channels.	1	[51]
CMS measurement of the t -channel single-top and anti-top cross sections. $\sigma_t \sigma_{\bar{t}} \sigma_{t+\bar{t}} R_t$.	1 1 1 1	[20]	CMS four-top search in the multi-lepton and same-sign dilepton channels.	1	[61]
ATLAS s -channel single-top cross section measurement.	1	[27]	CMS four-top search in the single-lepton and opposite-sign dilepton channels.	1	[59]
CMS tW cross section measurement.	1	[21]	CMS $t\bar{t}b\bar{b}$ cross section measurement in the all-jet channel.	1	[57]
ATLAS tW cross section measurement in the single lepton channel.	1	[41]	CMS $t\bar{t}b\bar{b}$ cross section measurement in the dilepton channel.	1	[64]
ATLAS tW cross section measurement in the dilepton channel.	1	[22]			

Fisher information breakdown

C_i	EWPO	LEP WW	Run 1 SS	Run 2 SS	STXS	LHC WW	WZ	Zjj	$t\bar{t}$	$W_{\text{hel.}}$	tX	$t\bar{t}V$
C_{HWB}	51	—	7	14	28	—	—	—	—	—	—	—
C_{HD}	100	—	—	—	—	—	—	—	—	—	—	—
C_{ll}	99	—	—	—	—	—	—	—	—	—	—	—
$C_{Hl}^{(3)}$	99	—	—	—	—	—	—	—	—	—	—	—
$C_{Hl}^{(1)}$	100	—	—	—	—	—	—	—	—	—	—	—
C_{He}	100	—	—	—	—	—	—	—	—	—	—	—
$C_{Hq}^{(3)}$	89	1	—	—	2	—	6	—	—	—	—	—
$C_{Hq}^{(1)}$	99	—	—	—	—	—	—	—	—	—	—	—
C_{Hd}	99	—	—	—	—	—	—	—	—	—	—	—
C_{Hu}	98	—	—	—	1	—	—	—	—	—	—	—
$C_{H\Box}$	—	—	22	46	32	—	—	—	—	—	—	—
C_{HG}	—	—	22	42	36	—	—	—	—	—	—	—
C_{HW}	—	—	14	29	56	—	—	—	—	—	—	—
C_{HB}	—	—	14	29	57	—	—	—	—	—	—	—
C_W	—	3	—	—	—	—	13	84	—	—	—	—
C_G	—	—	—	—	—	—	—	—	43	—	—	56
$C_{\tau H}$	—	—	22	45	34	—	—	—	—	—	—	—
$C_{\mu H}$	—	—	5	95	—	—	—	—	—	—	—	—
C_{bH}	—	—	19	35	47	—	—	—	—	—	—	—
C_{tH}	—	—	21	45	34	—	—	—	—	—	—	—
$C_{HQ}^{(3)}$	99	—	—	—	—	—	—	—	—	—	—	—
$C_{HQ}^{(1)}$	100	—	—	—	—	—	—	—	—	—	—	—
C_{Ht}	—	—	—	—	—	—	—	—	—	—	—	100
C_{tG}	—	—	13	29	24	—	—	—	24	—	—	9
C_{tW}	—	—	—	—	—	—	—	—	—	84	15	—
C_{tB}	—	—	—	—	—	—	—	—	—	—	—	100
$C_{Qq}^{3,1}$	—	—	—	—	—	—	—	—	—	—	100	—
$C_{Qq}^{3,8}$	—	—	—	—	—	—	—	—	87	—	—	13
$C_{Qq}^{1,8}$	—	—	—	—	—	—	—	—	82	—	—	17
C_{Qu}^8	—	—	—	—	—	—	—	—	91	—	—	7
C_{Qd}^8	—	—	—	2	—	—	—	—	92	—	—	6
C_{tq}^8	—	—	—	1	—	—	—	—	89	—	—	10
C_{tu}^8	—	—	—	—	—	—	—	—	96	—	—	3
C_{td}^8	—	—	—	2	—	—	—	—	92	—	—	5

Technical details

$$\mu_X \equiv \frac{X}{X_{SM}} = 1 + \sum_i a_i^X \frac{C_i}{\Lambda^2} + \mathcal{O}\left(\frac{1}{\Lambda^4}\right)$$

Exp. data: `HEPdata`, `WebPlotDigitizer`, ...

- Construct ‘signal strength’, w.r.t. SM prediction from exp. paper
- Otherwise computed with `MG5`, `fastnlo`, directly from theory papers
- Combine all sources of uncertainty in quadrature (stat., syst., th.)

Technical details

$$\mu_X \equiv \frac{X}{X_{SM}} = 1 + \sum_i a_i^X \frac{C_i}{\Lambda^2} + \mathcal{O}\left(\frac{1}{\Lambda^4}\right)$$

Exp. data: `HEPdata`, `WebPlotDigitizer`, ...

- Construct ‘signal strength’, w.r.t. SM prediction from exp. paper
- Otherwise computed with `MG5`, `fastnlo`, directly from theory papers
- Combine all sources of uncertainty in quadrature (stat., syst., th.)

Theory predictions: `MG5 (SMEFTsim & SMEFTatNLO)`

- LO, parton-level, linear dependence in (α, G_F, M_Z) scheme
- Tree-level + 1-loop gluon fusion Higgs production
- a_i : Effects from production, decays, total width
- No theory error from EFT, assume SM error dominant

The code

fitmaker <https://gitlab.com/kenmimasu/fitrepo>
public-friendly version w/ example notebooks in progress

Main analysis: linearised least-squares fit

$$\chi^2(C_i) = (\vec{y} - \vec{\mu}(C_i))^T \mathbf{V}^{-1} (\vec{y} - \vec{\mu}(C_i)) \quad \mu_\alpha(C_i) = \mu_\alpha^{\text{SM}} + \mathbf{H}_{\alpha i} C_i$$

The code

fitmaker <https://gitlab.com/kenmimasu/fitrepo>
public-friendly version w/ example notebooks in progress

Main analysis: linearised least-squares fit

$$\chi^2(C_i) = (\vec{y} - \vec{\mu}(C_i))^T \mathbf{V}^{-1} (\vec{y} - \vec{\mu}(C_i)) \quad \mu_\alpha(C_i) = \mu_\alpha^{\text{SM}} + \mathbf{H}_{\alpha i} C_i$$

Best fit $\hat{\vec{C}} = (\mathbf{H}^T \mathbf{V}^{-1} \mathbf{H})^{-1} \mathbf{H}^T \mathbf{V}^{-1} (\vec{y} - \vec{\mu}^{\text{SM}}) \equiv \mathbf{F}^{-1} \vec{\omega}$

$$\mathbf{F} \equiv \mathbf{H}^T \mathbf{V}^{-1} \mathbf{H} \quad , \quad \vec{\omega} \equiv \mathbf{H}^T \mathbf{V}^{-1} (\vec{y} - \vec{\mu}^{\text{SM}}) ,$$

Fisher information

The code

fitmaker <https://gitlab.com/kenmimasu/fitrepo>
public-friendly version w/ example notebooks in progress

Main analysis: linearised least-squares fit

$$\chi^2(C_i) = (\vec{y} - \vec{\mu}(C_i))^T \mathbf{V}^{-1} (\vec{y} - \vec{\mu}(C_i)) \quad \mu_\alpha(C_i) = \mu_\alpha^{\text{SM}} + \mathbf{H}_{\alpha i} C_i$$

Best fit $\hat{\vec{C}} = (\mathbf{H}^T \mathbf{V}^{-1} \mathbf{H})^{-1} \mathbf{H}^T \mathbf{V}^{-1} (\vec{y} - \vec{\mu}^{\text{SM}}) \equiv \mathbf{F}^{-1} \vec{\omega}$

$$\mathbf{F} \equiv \mathbf{H}^T \mathbf{V}^{-1} \mathbf{H} \quad , \quad \vec{\omega} \equiv \mathbf{H}^T \mathbf{V}^{-1} (\vec{y} - \vec{\mu}^{\text{SM}}) ,$$

Fisher information

$$\mathbf{F}^{-1} \equiv \mathbf{U} \quad \text{Covariance matrix of least-squares estimators}$$

The code

fitmaker <https://gitlab.com/kenmimasu/fitrepo>
public-friendly version w/ example notebooks in progress

Main analysis: linearised least-squares fit

$$\chi^2(C_i) = (\vec{y} - \vec{\mu}(C_i))^T \mathbf{V}^{-1} (\vec{y} - \vec{\mu}(C_i)) \quad \mu_\alpha(C_i) = \mu_\alpha^{\text{SM}} + \mathbf{H}_{\alpha i} C_i$$

Best fit $\hat{\vec{C}} = (\mathbf{H}^T \mathbf{V}^{-1} \mathbf{H})^{-1} \mathbf{H}^T \mathbf{V}^{-1} (\vec{y} - \vec{\mu}^{\text{SM}}) \equiv \mathbf{F}^{-1} \vec{\omega}$

$$\mathbf{F} \equiv \mathbf{H}^T \mathbf{V}^{-1} \mathbf{H} \quad , \quad \vec{\omega} \equiv \mathbf{H}^T \mathbf{V}^{-1} (\vec{y} - \vec{\mu}^{\text{SM}}) ,$$

Fisher information

$\mathbf{F}^{-1} \equiv \mathbf{U}$ Covariance matrix of least-squares estimators

$(\chi_{SM}^2, \hat{\vec{C}}, \mathbf{U})$ fully characterise likelihood

- Individual, profiled/marginalised bounds & correlations
- Principal component analysis (eigensystem of F)

Implemented as part
of the **fitmaker**
framework

The code

fitmaker <https://gitlab.com/kenmimasu/fitrepo>
public-friendly version w/ example notebooks in progress

Main analysis: linearised least-squares fit

$$\chi^2(C_i) = (\vec{y} - \vec{\mu}(C_i))^T \mathbf{V}^{-1} (\vec{y} - \vec{\mu}(C_i)) \quad \mu_\alpha(C_i) = \mu_\alpha^{\text{SM}} + \mathbf{H}_{\alpha i} C_i$$

Best fit $\hat{\vec{C}} = (\mathbf{H}^T \mathbf{V}^{-1} \mathbf{H})^{-1} \mathbf{H}^T \mathbf{V}^{-1} (\vec{y} - \vec{\mu}^{\text{SM}}) \equiv \mathbf{F}^{-1} \vec{\omega}$

$$\mathbf{F} \equiv \mathbf{H}^T \mathbf{V}^{-1} \mathbf{H} \quad , \quad \vec{\omega} \equiv \mathbf{H}^T \mathbf{V}^{-1} (\vec{y} - \vec{\mu}^{\text{SM}}) ,$$

Fisher information

$\mathbf{F}^{-1} \equiv \mathbf{U}$ Covariance matrix of least-squares estimators

$(\chi_{SM}^2, \hat{\vec{C}}, \mathbf{U})$ fully characterise likelihood

- Individual, profiled/marginalised bounds & correlations
- Principal component analysis (eigensystem of F)

Implemented as part
of the **fitmaker**
framework

Also nested sampling routine for general likelihoods

The code

fitmaker <https://gitlab.com/kenmimasu/fitrepo>
public-friendly version w/ example notebooks in progress

Database of input measurements encoded in **.json** format

- Values, errors, metadata,...

The code

fitmaker <https://gitlab.com/kenmimasu/fitrepo>
public-friendly version w/ example notebooks in progress

Database of input measurements encoded in `.json` format

- Values, errors, metadata,...

Python-class based definition of theoretical models

- Predictions for observables can be hard-coded
- ...or read-in from a `.json` file

The code

fitmaker <https://gitlab.com/kenmimasu/fitrepo>
public-friendly version w/ example notebooks in progress

Database of input measurements encoded in `.json` format

- Values, errors, metadata,...

Python-class based definition of theoretical models

- Predictions for observables can be hard-coded
- ...or read-in from a `.json` file

$$\mu_{H \rightarrow 4\ell}^{ggF} = 0.98^{+0.12}_{-0.11}$$

```
{
  "observable_name": "mu_ggF_H_ZZ_13",
  "measurement_name": "mu_ggF_H_ZZ_CMS_Run2",
  "CDS": "http://cds.cern.ch/record/2706103",
  "reportnumber": "CMS-PAS-HIG-19-005",
  "DOI": "",
  "date": "2020/01/10",
  "experiment": "CERN LHC Run 2, CMS",
  "description": "Higgs boson signal strength for",
  "value": 0.98,
  "uncertainty": {
    "tot": [0.12, 0.11]
  },
  "uncertainty_sigma": 1,
  "th_flat": true
}
```

$$\mu^{ggF} = 1 + 35.8C_{HG} - 0.122C_{tH} - 0.959C_{tG} - 0.121C_{H\Box} + \dots$$

```
{
  "observable": "ggF0j",
  "params": ["CHG", "CuH", "CuG", "CHbox"],
  "constant": 1.0,
  "linear": [35.8, -0.122, 0.959, -0.121],
  "quadratic": [
    [321.0, -1.095, 8.45, -1.085],
    [-1.095, 0.00371, -0.02925, 0.003695],
    [8.45, -0.02925, 0.23, -0.0291],
    [-1.085, 0.003695, -0.0291, 0.00367]
  ],
  "lambda_gen": 1000.0
}
```

SMEFT@NLO

Loops & SMEFT: active field in recent years

- Non-universal K-factors in EFT space \Leftrightarrow new information at NLO
- Loop-induced sensitivity (e.g. $gg \rightarrow H$)
- Control theoretical uncertainties
- Experimental interest in higher precision for SMEFT analyses/interpretations

SMEFT@NLO

Loops & SMEFT: active field in recent years

- Non-universal K-factors in EFT space \Leftrightarrow new information at NLO
- Loop-induced sensitivity (e.g. $gg \rightarrow H$)
- Control theoretical uncertainties
- Experimental interest in higher precision for SMEFT analyses/interpretations

Challenge: many processes x many operators

- LO \Rightarrow NLO = more cross-talk/operators/complexity
- Automated tools for fixed-order/NLO+PS are essential to the LHC programme

SMEFT@NLO

Loops & SMEFT: active field in recent years

- Non-universal K-factors in EFT space \Leftrightarrow new information at NLO
- Loop-induced sensitivity (e.g. $gg \rightarrow H$)
- Control theoretical uncertainties
- Experimental interest in higher precision for SMEFT analyses/interpretations

Challenge: many processes x many operators

- LO \Rightarrow NLO = more cross-talk/operators/complexity
- Automated tools for fixed-order/NLO+PS are essential to the LHC programme


Solution: SMEFT@NLO


- UFO model for MadGraph5_aMC@NLO
- Process-independent implementation: SMEFT in top-specific flavor limit

SMEFT@NLO in STXS

Gluon fusion Simplified Template Cross Section bins


- LO in the SM is one-loop
- Tree-EFT x loop-SM + loop-EFT x loop-SM interference terms
- Heavy top limit is OK for 0-jet, breaks down at high- p_T





Céline Degrande, Gauthier Durieux, Fabio Maltoni, Ken Mimasu, Eleni Vryonidou & Cen Zhang, [arXiv:2008.11743](#)

The implementation is based on the Warsaw basis of dimension-six SMEFT operators, after canonical normalization. Electroweak input parameters are taken to be G_F , M_Z , M_W . The CKM matrix is approximated as a unit matrix, and a $U(2)_q \times U(2)_u \times U(3)_d \times (U(1)_l \times U(1)_e)^3$ flavor symmetry is enforced. It forbids all fermion masses and Yukawa couplings except that only of the top quark. The model therefore implements the five-flavor scheme for PDFs.

A new coupling order, `NP=2`, is assigned to SMEFT interactions. The cutoff scale `Lambda` takes a default value of 1 TeV^{-2} and can be modified along with the Wilson coefficients in the `param_card`. Operators definitions, normalisations and coefficient names in the UFO model are specified in [definitions.pdf](#) . The notations and normalizations of top-quark operator coefficients comply with the LHC TOP WG standards of [1802.07237](#). Note however that the flavor symmetry enforced here is slightly more restrictive than the baseline assumption there (see the [dim6top page](#) for more information). This model has been validated at tree level against the `dim6top` implementation (see [1906.12310](#) and the [comparison details](#)).

Current implementation

UFO model: [SMEFTatNLO_v1.0.tar.gz](#) 

The current implementation imposes CP conservation. In the quark sector, it focuses primarily on top-quark interactions. The light-quark current operator, qqHDH, uuHDH, ddHDH, with coefficients `cpq3i`, `cpqMi`, `cpu`, `cpd` are however included. The triple-gluon operator, with coefficient `cG`, is currently not available (see the loop-capable `GGG` implementation). Vertices including more than four scalars or four leptons are not included. Scalar and tensor `QQ11` operators, with coefficients `ct1S3`, `ct1T3`, and `cb1S3`, break our flavor symmetry assumption and are not available for one-loop computations. Top-quark flavor-changing interactions, not compatible with the imposed flavor symmetry, are not included (see the loop-capable [TopFCNC](#) implementation).

Unlike prescribed by the LHC TOP WG, the top quark chromomagnetic-dipole operator coefficient `ctG` is normalized with a factor of the strong coupling, g_s . This normalization factor temporarily ensures compatibility with the 2.X.X series of MadGraph5_aMC@NLO but may be dropped in the future. As with every other appearance of this coupling in MadGraph5_aMC@NLO, its value is renormalisation-group evolved to the QCD renormalization scale (set in the `run_card`).

```
MG5_aMC>import model SMEFTatNLO
```


```
MG5_aMC>generate p p > t t~ NP=2 [QCD]
```

```
MG5_aMC>output
```



```
MG5_aMC>launch
```

Top couplings via Higgs

Tree



Loop



- Significant effects on individual observable sensitivities, e.g., tZ
- Loop-induced processes bring new constraints ($h \rightarrow \gamma\gamma/\gamma Z$)