



# Probing $\theta_{13}$ With The Daya Bay Antineutrino Detectors


Bryce Littlejohn, on behalf of the  
Daya Bay Collaboration

5/11/10





- Aims to measure reactor  $\bar{\nu}_e$  disappearance:



- 6 reactor cores, 3 sites
- 4 near detectors, 2 sites,  $\sim 900$  evts/day/det
- 4 far detectors, 1 site,  $\sim 90$  evts/day/det
- Significant overburden
- RPC, Muon water veto

	DYB Site	LA Site	Far Site
Depth (m rock)	98	112	350
Bkg/Sig	0.3%	0.2%	0.2%




# How does Daya Bay probe $\theta_{13}$ ?

- Detect short-baseline reactor antineutrino disappearance:

$$P_{13} \approx 1 - \sin^2 2\theta_{13} \sin^2 \left[ 1.27 \Delta m_{13}^2 (eV^2) \frac{L(km)}{E_\nu (GeV)} \right]$$



- A “clean” measurement of  $\theta_{13}$





Near Sites

Far Site

- 0.1% Gd-doped liquid scintillator as an inverse-beta target:





$$E_{e^+} = E_{\bar{\nu}_e} - 1.29\text{MeV}$$



Daya Bay Monte Carlo Data




- 8 'identical' 3-zone detectors:





Target Mass: 20 tons  
Energy Resolution:  $12\%/\sqrt{E}$   
Light Yield:  $\sim 120$  photoelectrons/MeV



- First 2 ADs being assembled:
  - SSV, bottom reflector: Installed, surveyed
  - Outer acrylic vessels installed, 1 inner vessel installed
  - PMT ladders being populated, installed late May, June
- Next few months:
  - Close stainless steel lids: June, July
  - Attach calibration boxes and overflow tanks: June, July
  - Dry-run data taking in July






May 11, 2010







# AD Liquids: Description and Status



Daya Bay LS Hall



# Measurement Method



$$\frac{N_f}{N_n} = \left( \frac{N_{p,f}}{N_{p,n}} \right) \left( \frac{L_n}{L_f} \right)^2 \left( \frac{\epsilon_f}{\epsilon_n} \right) \left[ \frac{P_{\text{sur}}(E, L_f)}{P_{\text{sur}}(E, L_n)} \right]$$





- Daya Bay is a systematics-limited experiment
  - With near-far ratio measurement, detector systematics become dominant:

Source of uncertainty		Baseline	Goal
Number of Protons		0.3%	0.1%
Detector Efficiency	Energy Cuts	0.2%	0.1%
	Time Cuts	0.1%	0.03%
	H/Gd Ratio	0.1%	0.1%
	n multiplicity	0.05%	0.05%
	Trigger	0.01%	0.01%
	Live Time	<0.01%	<0.01%
	Total Efficiency	0.25%	0.15%
Total detector uncorrelated uncertainty		0.38%	0.18%

- Largest: number of protons and 6MeV energy cut efficiency
- Working hard to minimize these dominant uncertainties


200-ton Gd-LS reservoir



20-ton ISO tank

ISO Gd-LS weighing tank

filling platform with clean room




load cell accuracy < 0.02%



Coriolis mass flowmeters < 0.1%  
For food, beverage, pharmaceutical and chemical applications!

- To beat systematics goal:

- Redundant mass measurement systems
  - Measure mass to 0.02%: +/- 4 kg out of 20 tons
- Filling detectors in pairs from a common GdLS reservoir
  - Reduce differences between detector liquids




filling "pairs" of detectors





- In practice, detectors will not be physically identical
  - Real question: are they identical enough to have similar physics response?
- Characterize as-built detectors - identify differences

Variations  
In Acrylic  
Vessel  
Transmittance





Variations  
In acrylic vessel  
thickness, i.e.  
non-scintillating  
(dead) volume

- Use MC simulation to see if differences affect physics performance



- Spring 2011: Near site turn-on
  - Reactor flux studies
  - Comparison of near site detectors and evaluation of systematics
- Spring 2012: Far site turn-on
  - Should achieve our goal sensitivity of  $\sin^2 2\theta_{13} < 0.01$  at 90% CL





- AD 1,2 assembly is well underway; done by Summer 2010
- R&D has shown that we can meet or exceed goal systematics
  - Detector construction and filling in pairs
  - Mass Measurement R&D
- Spring 2011: Data taking at the near hall will start
- Spring 2012: Full data taking start
- Can reach  $\sin^2 2\theta_{13} < 0.01$
- Chris White will provide more details next
- Questions?



- Thanks!



- Current limits on  $\theta_{13}$  set by:
  - MINOS: accelerator  $\nu_e$  appearance
  - CHOOZ: reactor  $\bar{\nu}_e$  disappearance

