


Trilepton Production at LHC

Standard model sources and beyond



Zack Sullivan

Southern Methodist University


April 28, 2008

Based on Z.S., E. Berger, ANL-HEP-PR-08-21, hep-ph:0805.xxxx

Outline

1. Dileptons and trileptons at LHC
2. How heavy flavors (b, c) yield isolated leptons
3. SUSY signal vs. leptons from heavy flavors
4. Improved cuts
5. Conclusions

Motivation: Dileptons at LHC and b quark decays




Higgs decays to W^+W^- to dileptons is expected to give the largest significance for $135 < M_H < 219$ GeV at LHC. ATLAS TDR V.2

A study of heavy-flavor (b, c) decay to leptons found $b\bar{b} + Wb\bar{b} + Wc + \text{single-top} + \dots$ is $> 50 \times$ the direct WW background.

Conclusion: Isolation does not remove leptons from heavy flavor decays!

Z.S., E. Berger, PRD74, 033008 (06)

Motivation: Dileptons at LHC and b quark decays




Higgs decays to W^+W^- to dileptons is expected to give the largest significance for $135 < M_H < 219$ GeV at LHC. ATLAS TDR V.2

A study of heavy-flavor (b, c) decay to leptons found $b\bar{b} + Wb\bar{b} + Wc + \text{single-top} + \dots$ is $> 50\times$ the direct WW background.


Conclusion: Isolation does not remove leptons from heavy flavor decays!

Z.S., E. Berger, PRD74, 033008 (06)



Solution: The second lepton p_T falls exponentially.

So raise the cut: $p_{Tl_2} > 10$ GeV $\Rightarrow p_{Tl_2} > 20$ GeV.




Leading edge 20 GeV lower!

$H \rightarrow WW$ survives!

$b\bar{b} \rightarrow b\bar{b}/30, W+X \rightarrow W+X/10,$
 $t+X \rightarrow t+X/5$


Motivation: Trileptons at LHC



$\tilde{\chi}_1^\pm \tilde{\chi}_2^0 \rightarrow l^+ l^- l^\pm + \cancel{E}_T$ is a golden signature of supersymmetry.

CMS and ATLAS both have analyses designed to observe this signal.


CMS TDR V.2&Note 2006/113; ATLAS CSC 7



WZ was expected to be the largest source of low- p_T trileptons at LHC.

$W\gamma^*$ has not previously been included.


Motivation: Trileptons at LHC



$\tilde{\chi}_1^\pm \tilde{\chi}_2^0 \rightarrow l^+ l^- l^\pm + \cancel{E}_T$ is a golden signature of supersymmetry.


CMS and ATLAS both have analyses designed to observe this signal.

CMS TDR V.2&Note 2006/113; ATLAS CSC 7



WZ was expected to be the largest source of low- p_T trileptons at LHC.

$W\gamma^*$ has not previously been included.




How important are leptons from heavy flavor (b, c) decays?

There are MANY potential processes:


$bZ/\gamma, b\bar{b}Z/\gamma, cZ/\gamma, c\bar{c}Z/\gamma, b\bar{b}W, c\bar{c}W, t\bar{t}, tW, t\bar{b}$

NOTE: All photons are virtual, and split to $l^+ l^-$

Physics of isolated leptons from b decay



Physics of isolated leptons from b decay



Prob. isolated muon



= Prob. producing muon
× Prob. B remnants missed

- Muons that pass isolation take nearly all p_T
- ~Nearly all isolated muons point back to primary vertex.

C. Wolfe, CDF internal

- Isolation leaves $\sim 7.5 \times 10^{-3} \mu/b$
 $\gg 10^{-4}$ per light jet

Physics of isolated leptons from b decay



Prob. isolated muon

= Prob. producing muon
× Prob. B remnants missed

- Muons that pass isolation take nearly all p_T
- ~Nearly all isolated muons point back to primary vertex.

C. Wolfe, CDF internal


- Isolation leaves $\sim 7.5 \times 10^{-3} \mu/b$
 $\gg 10^{-4}$ per light jet

Harder b 's can give isolated e 's, because e cuts must allow more energy in the calorimeter

It is difficult to reduce this without losing efficiency for primary e .

Isolation is not extremely effective for leptons from b decay.

Isolated leptons from b/c production & decay



Fold in $b\bar{b}$ production.


- A large fraction of events with $b \rightarrow \mu/e$ have isolated μ/e .

More isolated e than μ per b .

- 1/2 of all isolated μ come from b with $p_{Tb} < 20$ GeV.

It is common for analyses to start simulations with $p_{Tb} > 20$ GeV.


Isolated leptons from b/c production & decay



Fold in $b\bar{b}$ production.

- A large fraction of events with $b \rightarrow \mu/e$ have isolated μ/e .
More isolated e than μ per b .
- 1/2 of all isolated μ come from b with $p_{Tb} < 20$ GeV.

It is common for analyses to start simulations with $p_{Tb} > 20$ GeV.



Fold in $c\bar{c}$ production.

The story repeats for c decays

1 twist: D decays have many pions

π^\pm fake e at $\sim 10^{-4}$


\Rightarrow Large " e_{iso} " rate

Trileptons: SUSY & SM at CMS w/ 30 fb^{-1}

Channel	$N^l = 3,$ NoJets	M_{ll}^{OSSF} < 75 GeV
LM9	248	243
LM7	126	123
LM1	46	44
WZ/γ	1880	538
$t\bar{t}$	1540	814
tW	273	146
$t\bar{b}$	1.1	1.0
bZ/γ	14000	6870
cZ/γ	3450	1400
$b\bar{b}Z/\gamma$	8990	2220
$c\bar{c}Z/\gamma$	4680	1830
$b\bar{b}W$	9.1	7.6
$c\bar{c}W$	0.19	0.15

Analysis cuts:

- 3 leptons
- No jets ($E_{Tj} > 30 \text{ GeV}$)
- Remove Z peak
(demand $M_{ll}^{\text{OSSF}} < 75 \text{ GeV}$)




Z+heavy flavor decays are
10× $WZ/\gamma + t\bar{t}$!

Two additional cuts: \cancel{E}_T and angular correlations

Leptons from SUSY decays are SOFT \Rightarrow Cannot raise p_{Tl} cut.

Missing E_T



Z/γ +heavy flavors – no intrinsic \cancel{E}_T

Comes from misreconstruction,
energy lost down beam pipe

Natural \cancel{E}_T in SUSY points low as well

$\tilde{\chi}_1^0$'s partially balance out


A \cancel{E}_T cut demanding

$\cancel{E}_T > 30\text{--}40$ GeV is very effective

Two additional cuts: \cancel{E}_T and angular correlations

Leptons from SUSY decays are SOFT \Rightarrow Cannot raise p_{Tl} cut.

Missing E_T




Z/γ +heavy flavors – no intrinsic \cancel{E}_T
Comes from misreconstruction,
energy lost down beam pipe

Natural \cancel{E}_T in SUSY points low as well
 $\tilde{\chi}_1^0$'s partially balance out

A \cancel{E}_T cut demanding
 $\cancel{E}_T > 30\text{--}40$ GeV is very effective

\cancel{E}_T is poorly measured

Angular correlations



Angles measured extremely well
All combinations different (θ_{12}^{CM} shown)

Demand $\theta_{12}^{\text{CM}} > 45^\circ$, $\theta_{13}^{\text{CM}} > 40^\circ$,
 $\theta_{23}^{\text{CM}} < 160^\circ$

Reduces B by 30% for 5% loss of S
Not optimized

Trileptons: SUSY & SM at CMS (+new cuts)

Channel	$N^l = 3,$ NoJets	M_{ll}^{OSSF} < 75 GeV	$\cancel{E}_T > 30$ GeV	Angular cuts
LM9	248	243	160	150
LM7	126	123	89	85
LM1	46	44	33	32
WZ/γ	1880	538	325	302
$t\bar{t}$	1540	814	696	672
tW	273	146	123	121
$t\bar{b}$	1.1	1.0	0.77	0.73
bZ/γ	14000	6870	270	177
cZ/γ	3450	1400	45	35
$b\bar{b}Z/\gamma$	8990	2220	119	103
$c\bar{c}Z/\gamma$	4680	1830	69	35
$b\bar{b}W$	9.1	7.6	5.6	5.3
$c\bar{c}W$	0.19	0.15	0.12	0.11

Pure QCD background to trileptons

CMS estimates $jjj \rightarrow lll < 5$ events in 30 fb^{-1}

What about $b\bar{b}b\bar{b}$, $b\bar{b}c\bar{c}$, $c\bar{c}c\bar{c}$?

We cannot simulate this directly in our lifetimes ($\sim 10^3$ CPU years)

Estimate 3 sources of $b\bar{b}b\bar{b}$ for 30 fb^{-1}

1. Direct $b\bar{b}b\bar{b}$: ~ 500 events

Use $Wb\bar{b}$ to estimate $P(b \rightarrow \mu_{\text{iso}})$: $\sigma_{b\bar{b}b\bar{b}} \times (7.5 \times 10^{-3})^3$

2. Multiple interactions: ~ 600 events

10 interactions $\times \sigma_{b\bar{b}}^2 / \sigma_{\text{inelastic}}^{\text{Tot}}$

3. Multiple scattering, gluon splitting: $\sim 10^3$ events


Note that K factors could be as high as 5.5

A. Del Fabbro, D. Treleani, PRD66, 074012 (02)

Scaling results from Z.S., E.L. Berger, PRD 74, 033008 (06),
the \cancel{E}_T cut should remove nearly all of these.

Importance of the virtual photon



Simulations of WZ based on PYTHIA do not include virtual photons.



Nearly 1/2 of the trilepton background from WZ/γ is from $W\gamma^*$ alone.

Matrix elements that include virtual photons are important when observing low- p_T leptons.

(p_{Tl} spectra after M_{ll}^{OSSF} cut)



Significance of SUSY point LM9 in 30 fb^{-1}

1. **Our calculations are LO.**
NLO K -factors are large (1.5–2) on most processes,
BUT, jet veto will reduce this.
2. **ISR is not well determined**
The rate of $> 30 \text{ GeV}$ jets can be changed by a factor of **4**
depending on assumptions in PYTHIA about ISR.

We present our calculation, and one that scales down B by 4 to show the range of possible significances

	$N^l = 3,$ NoJets	M_{ll}^{OSSF} < 75 GeV	$\cancel{E}_T > 30 \text{ GeV}$	Angular cuts
S/\sqrt{B}_{LM9}	1.33	2.07(1.79)	3.93(3.74)	3.94(3.79)
$S/\sqrt{B}_{\text{LM9}}^{\text{CMS } j}$	2.63	4.09(3.54)	7.78(7.39)	7.79(7.49)

(Parentheses include leptons from fakes from CMS Table 6, Note 2006/113)

We will not know which ISR estimate is correct until we measure it at LHC

Conclusions

(Z.S., E. Berger, ANL-HEP-PR-08-21, hep-ph:0805.xxxx)


1. Heavy-flavor (b, c) decays to leptons dominate low- p_T isolated leptons at LHC
Trileptons from $Z/\gamma^* + \text{heavy flavors (HF)} \sim 10\times$ all other backgrounds
2. When modeling low- p_T leptons, virtual photons cannot be ignored
 $WZ/\gamma^* \sim 1.7 \times WZ$ after cuts
3. Raising minimum p_T is not viable for SUSY signal, but other cuts work:
 - (a) Require $\cancel{E}_T > 30$ GeV, $Z/\gamma^* + \text{HF} \rightarrow Z/\gamma^* + \text{HF}/30$ Hard to measure low \cancel{E}_T
 - (b) Impose cuts on well-measured angles, $Z/\gamma^* + \text{HF}$ reduced by 30%
4. Overall normalization is dominated by assumptions regarding ISR
Huge uncertainties in effectiveness of jet veto
If large ISR exists, may want to loosen jet veto to recover SUSY signal
ISR questions will be resolved with initial data from LHC

Any signal that has low- p_T leptons MUST consider the background from heavy flavor (b, c) decays


BACKUPS

Other angular correlations

Angles are well-measured, and defined in the trilepton CM frame.



Suggested cut: $\theta_{13}^{\text{CM}} > 40^\circ$




Suggested cut: $\theta_{23}^{\text{CM}} < 160^\circ$

These cuts are almost free, and not optimized.


5% signal decrease, but 30% background decrease

CMS SUSY points LM1, LM7

Representative opposite-sign same-flavor (OSSF) invariant masses



Signal endpoint above Z -peak cut
and signal is small



LM7 similar to LM9, but smaller