

Exploring the Phenomenology of the Noncommutative Standard Model with WW Scattering

J.A. Conley J.L. Hewett

Theory Group
Stanford Linear Accelerator Center

PHENO Symposium, April 28-30, 2008

arXiv:0805.????

Outline.

- 1 Noncommutative field theory.
- 2 Results.
 - Unitarity.
 - Phenomenology.

The Long History of Noncommutative Theories.

- Noncommutative (NC) physics first proposed to solve divergence problems in QFT. (Heisenberg; Pauli; Snyder)
- NC geometry became its own field of mathematics, and was used to “derive” SM. (Connes)
- NCQFT shown to arise from low energy limit of string theory with a B field. (Seiberg, Witten)
- NC gauge theories are formulated; two versions of NCSM are developed. (Wess *et. al.*; Chaichian *et. al.*)

The Basics of Noncommutativity.

Fundamental assumption.

- $[x_\mu, x_\nu] = i\theta_{\mu\nu} \equiv i\frac{c_{\mu\nu}}{\Lambda^2}$. Λ is “NC scale.”
- $c_{ij} \neq 0$ for $i, j \in \{1, 2, 3\}$ is called “space-space noncommutativity”.
- $c_{0i} \neq 0$ is called “space-time noncommutativity.”

Subtleties. (Things I won't talk about.)

- For space-time NC, perturbation theory must be formulated carefully—see time-ordered perturbation theory (Chaichian *et. al.*).
- So-called UV-IR mixing makes renormalization tricky, but possible (Filk; Minwalla *et. al.*; Petriello; Wulkenhaar; Buric *et. al.*; Martin, *et. al.*; ...).

Expressing Noncommutative Fields in Terms of Commutative Fields.

Moyal Star Product.

- For $U(n)$ gauge groups we can simply replace products with Moyal Star Products.
- $\hat{f}(\hat{x})\hat{g}(\hat{y}) \rightarrow f(x) \star g(y) = f(x) \exp \left\{ \frac{i}{2} \theta^{\mu\nu} \overleftarrow{\partial}_\mu \overrightarrow{\partial}_\nu \right\} g(y)$.

Seiberg-Witten Map (SWM).

- $SU(n)$ gauge groups don't close under \star -product.
- Can, however, impose consistency conditions on NC and normal gauge transformations.
- Doing so gives SWM, expanding NC quantities in terms of commuting quantities, order by order in $\theta_{\mu\nu}$.

Noncommutative Standard Model (NCSM).

Wess *et. al.* hep-ph/0111115, ...

Strategy.


- Take SM Lagrangian, and replace all fields and coordinates with NC counterparts.
- Using \star -product and SWM, expand \mathcal{L} in terms of commuting quantities, order-by-order in $\theta_{\mu\nu}$.

Result.

- No new particles (besides SM ones) appear in theory.
- At each order in θ , get many new interaction terms.
- Ambiguity in SWM for gauge kinetic terms gives three new free parameters $\kappa_1, \kappa_2, \kappa_3$.

All our calculations will be to first order in θ .

Feynman Rules.



$$\begin{aligned}
 &= -\frac{em_W^2}{2} f_{\mu\nu\rho}^A(p) + \frac{em_Z^2}{4} f_{\mu\nu\rho}^Z(p, q, r) \\
 &+ 2e \sin 2\theta_W K_{WWZ} \Theta_{\mu\nu\rho}(p, q, r), \text{ where}
 \end{aligned}$$

$$f_{\mu\nu\rho}^A(p) \equiv \theta_{\mu\nu} p_\rho + \theta_{\mu\rho} p_\nu + g_{\mu\nu}(\theta \cdot p)_\rho - g_{\nu\rho}(\theta \cdot p)_\mu + g_{\rho\mu}(\theta \cdot p)_\nu,$$

$$f_{\mu\nu\rho}^Z(p, q, r) \equiv \theta_{\mu\nu}(p - q)_\rho + \theta_{\nu\rho}(q - r)_\mu + \theta_{\rho\mu}(r - p)_\nu$$

$$- 2g_{\mu\nu}(\theta \cdot r)_\rho - 2g_{\nu\rho}(\theta \cdot p)_\mu - 2g_{\rho\mu}(\theta \cdot q)_\nu,$$

$$\Theta_{\mu\nu\rho}(p, q, r) \equiv \theta_{\mu\nu}(p \cdot r q_\rho - q \cdot r p_\rho) + (\theta \cdot p)_\mu(q \cdot r g_{\nu\rho} - q_\rho r_\nu)$$

$$- (\theta \cdot p)_\nu(q \cdot r g_{\rho\mu} - q_\rho r_\mu) - (\theta \cdot p)_\rho(q \cdot r g_{\mu\nu} - q_\mu r_\nu)$$

$$+ p \times q (r_\mu g_{\nu\rho} - r_\nu g_{\rho\mu}) + (\text{cyclic perms. of } \{p, q, r\} \text{ and } \{\mu, \nu, \rho\}),$$

$$\text{and } K_{WWZ} \equiv g^2 \kappa_2 / 2c_W^2.$$

Partial-Wave Unitarity in WW scattering.

Partial-Wave Unitarity.

- Decompose a scattering amplitudes as


$$A = 16\pi \sum_{l=0}^{\infty} a_l (2l+1) P_l(\cos \theta).$$
- The partial waves must obey

$$\text{Re}(a_l) \leq 1/2, \quad 0 \leq \text{Im}(a_l) \leq 1, \quad \text{and} \quad |a_l|^2 \leq 1.$$

$W^+ W^- \rightarrow W^+ W^-$ and $e^+ e^- \rightarrow W^+ W^-$ in NCSM.

- Amplitudes depend on $\theta_{\mu\nu}$ and κ_2 .
- Use partial-wave unitarity conditions on amplitudes to bound these parameters.
- Amplitudes are ϕ -dependent; maximize wrt ϕ to get “worst-case scenario” and apply conditions above.
- $e_L^- e_R^+ \rightarrow W_0^+ W_0^-$ gives strongest bound.


Unitarity bounds on NCSM.



WW scattering at colliders.


- Helicity-summed observables $\propto (c_{01} \sin \phi - c_{02} \cos \phi) / \Lambda^2$.
 - At LEP-II we look at $d\sigma/d \cos \theta d\phi$.
 - At LHC we look $d\sigma/dm_{WW}d\phi$.
- At ILC, distinguishing W helicities can give sensitivity to other parameters.
 - We use $d\sigma/d \cos \theta d\phi$ and $dA_{LR}/d \cos \theta d\phi$ to get search reach in $\kappa_2 - \Lambda$ plane.
 - We look at $d\sigma/d \cos \theta d\phi$ for different W helicities to determine $c_{\mu\nu}$, Λ , and κ_2 .

LEP-II search reach.




Observables summed
 over all final state
 helicities.

LHC search reach.




Using $\mathcal{L} = 100 \text{ fb}^{-1}$,
 and multiplying by
 branching ratio to
 semileptonic final
 states. Observables
 summed over all W
 helicities.

500 GeV ILC search reach.



Using $\mathcal{L} = 500 \text{ fb}^{-1}$
 and $P_{e^-} = 0.9$,
 $P_{e^+} = 0.6$,
 $\Delta P/P = 0.25\%$.
 Observables summed
 over all final state
 helicities.

1 TeV ILC search reach.



Using $\mathcal{L} = 500 \text{ fb}^{-1}$
 and $P_{e^-} = 0.9$,
 $P_{e^+} = 0.6$,
 $\Delta P/P = 0.25\%$.
 Observables summed
 over all final state
 helicities.

Comparison to Other Studies


- OPAL collaboration hep-ex/0303035: $e^+e^- \rightarrow \gamma\gamma$ gives $\Lambda > 141$ GeV.
- Alboteanu, Ohl, and Rückl, hep-ph/0608155: $pp \rightarrow Z\gamma$ gives search reach to $\Lambda \sim 1$ TeV.
- Alboteanu, Ohl, and Rückl, arXiv:0708.2359:

$(K_{Z\gamma\gamma}, K_{ZZ\gamma})$	$ \vec{E} ^2 = 1, \vec{B} = 0$	$\vec{E} = 0, \vec{B} ^2 = 1$
$K_0 \equiv (0, 0)$	$\Lambda \gtrsim 2$ TeV	$\Lambda \gtrsim 0.4$ TeV
$K_1 \equiv (-0.333, 0.035)$	$\Lambda \gtrsim 5.9$ TeV	$\Lambda \gtrsim 0.9$ TeV
$K_5 \equiv (0.095, 0.155)$	$\Lambda \gtrsim 2.6$ TeV	$\Lambda \gtrsim 0.25$ TeV
$K_3 \equiv (-0.254, -0.048)$	$\Lambda \gtrsim 5.4$ TeV	$\Lambda \gtrsim 0.9$ TeV


Table: Bounds on Λ from $e^+e^- \rightarrow Z\gamma$ at the ILC, for the minimal (first row) and nonminimal NCSM

Collider Observables.

SM vs. NCSM event rate for LHC, 100 fb^{-1}
 $\Lambda=600 \text{ GeV}$, $c_{01}=c_{02}=1$, $\kappa_2=3$, $m_{\text{WW}} \in [1.4, 1.6] \text{ TeV}$




SM vs. NCSM event rate for 1 TeV ILC, 500 fb^{-1}
 $\Lambda=2 \text{ TeV}$, $c_{01}=c_{02}=1$, $\kappa_2=1$




Polarized W 's at 1 TeV ILC.

$$W^-_0 W^+ + \frac{dN}{d\cos\theta} \text{ for } \phi < \pi, \kappa_2 = 0$$



$$W^-_0 W^+ + \frac{dN}{d\cos\theta} \text{ for } \phi < \pi, \kappa_2 = 1$$



$$\mathcal{L} = 500 \text{ fb}^{-1}, \Lambda = 10 \text{ TeV.}$$

Summary.

- The NCSM is a model of noncommutativity that could be observed in future experiments.
- Unitarity and LEP constrain but don't rule out the NCSM.
- At the LHC, the search reach is not much better than the unitarity bound.
- The ILC can probe large regions of NCSM parameter space and is sensitive to all of the parameters through W polarization measurements.
- Outlook
 - Implement unitarity more appropriately for ϕ -dependent amplitudes (using Chaichian *et. al.* hep-th/0305243).
 - Go beyond tree level and beyond leading order in θ (Alboreanu *et. al.* 0707.3595).
 - More realistic phenomenology: generate events, include detector effects, *etc.* (Alboreanu *et. al.*)