

Minimal Little Higgs Model and Dark Matter

Yang Bai

Theoretical Physics Department, Fermilab

2008 Phenomenology Symposium

[[arXiv:0801:1662](https://arxiv.org/abs/0801.1662)]


April 28, 2008

Outline

- 1 Motivation
 - Little Hierarchy Problem
 - Little Higgs Model
 - Simplify Little Higgs Model
- 2 The Model
 - Bosonic Sector
 - Fermionic Sector: Z_2 broken
 - Fermionic Sector: Z_2 unbroken
- 3 Dark Matter
 - Dark Matter
 - Relic Abundance
 - Direct Detection


Radiative Corrections to the Higgs Boson Mass

- The mass of the Higgs field is not stable against radiative corrections:




$$-\frac{3\lambda_t^2}{8\pi^2} \Lambda^2$$

$$\Lambda = 10 \text{ TeV}: \sim (2 \text{ TeV})^2$$



$$\frac{3(3g^2 + g_y^2)}{64\pi^2} \Lambda^2$$

$$\sim (700 \text{ GeV})^2$$



$$\frac{\lambda^2}{16\pi^2} \Lambda^2$$

$$\sim (500 \text{ GeV})^2$$


- Little hierarchy problem [LEP paradox] [Barbieri and Strumia, 2000]:
 - The mass of Higgs boson is less than 250 GeV.
 - The cutoff Λ of relevant higher-dimensional operators must be greater than 5-10 TeV.

Little Higgs Model

[Arkani-Hamed, Cohen and Georgi, 2001]

- Identify the Higgs doublet as a **pseudo-Nambu-Goldstone boson** (PNGB) of a spontaneously broken global symmetry.
- **Collective Symmetry Breaking**: two or more couplings are needed to explicitly break the global symmetry.
- **Consequence**: only logarithmically divergent potentials of the Higgs doublet are generated at one-loop level. The weak scale can be protected up to 5-10 TeV.


- Cancellations in the gauge sector:



- Gauge symmetries in various little Higgs models [$SU(3)_c$ is not included]:
 - The minimal moose model: $SU(3) \times SU(2) \times U(1)$.
 - The littlest Higgs model: $[SU(2) \times U(1)]^2$.
 - The simplest little Higgs model: $SU(3) \times U(1)$.
- Predict Z' , W' ; t' and partners of other light quarks; extra scalars including triplets and singlets.

Simplify Little Higgs Model

- How about the most minimal extension of the standard model gauge group: $SU(2) \times U(1) \times U(1)$?



- What is the symmetry for this cancellation?

Observations

- The field content under the gauge symmetry:

	$SU(2)_w$	$U(1)_1$	$U(1)_2$
H	2	1/2	1/2
S	1	5/3	-5/3

- The kinetic terms of scalars:

$$|(\partial_\mu + ig t^a W_\mu^a + i \frac{g'}{2\sqrt{2}}(B_{1\mu} + B_{2\mu}))H|^2 + |(\partial_\mu + i \frac{5g'}{3\sqrt{2}}(B_{1\mu} - B_{2\mu}))S|^2$$

- A Z_2 interchanging symmetry: $g_1 = g_2 = \sqrt{2}g'$

g' is the gauge coupling of $U(1)_Y$; g is the gauge coupling of $SU(2)_w$.

- The Λ^2 contributions to scalar masses from gauge bosons are:

$$V_g = \frac{3\Lambda^2}{64\pi^2} \left[(3g^2 + g'^2)HH^\dagger + \frac{100}{9}g'^2 SS^\dagger \right] + \dots,$$

$$\approx \frac{25g'^2\Lambda^2}{48\pi^2} [HH^\dagger + SS^\dagger] \propto \phi\phi^\dagger \Rightarrow \text{Approximate } U(3) \text{ global symmetry}$$

$$\sin^2 \theta_w = g'^2 / (g^2 + g'^2) \approx 0.23$$

Nonlinear Realization

- Write H and S together as a triplet of $U(3)$: $\phi = (H, S)^T$
- $\langle \phi \rangle = (0, 0, f)^T$ from underlying dynamics

$$\text{global symmetry: } U(3) \rightarrow U(2)$$

$$\text{gauge symmetry: } SU(2)_w \times U(1)_1 \times U(1)_2 \rightarrow SU(2)_w \times U(1)_Y$$

- Below the cutoff $\Lambda \approx 4\pi f$, the EFT contains 9-4=5 GB's.
 - One is eaten by the massive neutral gauge boson: $B' \equiv (B_1 - B_2)/\sqrt{2}$
 - The other 4 become PNGB's and identified as the Higgs doublet: h

$$\phi^T = f \left(\frac{ih}{\langle h \rangle} \sin \frac{\langle h \rangle}{f}, \cos \frac{\langle h \rangle}{f} \right) = (ih, f - \frac{\langle h \rangle^2}{2f}) + \dots$$

- The field dependent masses of gauge bosons are:

$$M_W^2(h) = c_w^2 M_Z^2(h) = \frac{1}{2} g^2 f^2 \sin^2 \frac{\langle h \rangle}{f} \quad M_{B'}^2(h) = \frac{50}{9} g'^2 f^2 \cos^2 \frac{\langle h \rangle}{f}$$

- Calculate the one-loop effective potential

$$V_{CW} = \frac{3}{32\pi^2} \Lambda^2 \text{Tr}[M_g^2] - \frac{3}{64\pi^2} \text{Tr}[M_g^4 \log(\frac{\Lambda^2}{M_g^2} + \frac{3}{2})]$$

- The Higgs mass contributions from the gauge sector:

$$m_h^2|_g = \frac{3g'^2\Lambda^2}{32\pi^2} \left(\frac{27-118s_w^2}{9s_w^2} \right) + \frac{3M_{B'}^4}{32\pi^2 f^2} \left(\log \frac{\Lambda^2}{M_{B'}^2} + 1 \right)$$

- For s_w^2 around 0.23, the Λ^2 term is even smaller than $\log \Lambda$ term

(Approx. $U(3)$ symmetry)

$$m_h^2|_g \approx -(87 \text{ GeV})^2 + (116 \text{ GeV})^2,$$

for $f = 800 \text{ GeV}$, $\Lambda = 10 \text{ TeV}$, $s_w^2 = 0.23$.

Z_2 Broken Model

- The field content under the gauge symmetry:

	$SU(3)_c$	$SU(2)_w$	$U(1)_1$	$U(1)_2$
H	1	2	1/2	1/2
S	1	1	5/3	-5/3
q_L	3	2	1/6	1/6
t_R	3	1	2/3	2/3
b_R	3	1	-1/3	-1/3
ψ_L	3	1	7/3	-1
ψ_R	3	1	7/3	-1

Only a colored vector-like quark added; gauge anomalies are still cancelled.

- The Yukawa couplings in the top sector are:

$$\mathcal{L}_t = y_1 (\bar{q}_L, \bar{\psi}_L) \phi t_R + y_2 f \bar{\psi}_L \psi_R = y_1 (\bar{q}_L \tilde{H} + \bar{\psi}_L S) t_R + y_2 f \bar{\psi}_L \psi_R + h.c.$$

Z_2 symmetry is manifestly broken

Z_2 Broken Model

- Higgs boson masses from the top sector:

$$m_h^2|_t = -\frac{3}{8\pi^2} y_t^2 m_{t'}^2 (\log \frac{\Lambda^2}{m_{t'}^2} + 1)$$

No Λ^2 contribution: collective breaking mechanism protects it.

- Spontaneously electroweak symmetry breaking:

$$m_h^2 = m_h^2|_g + m_h^2|_t < 0$$

Minimizing the full potential, we get a light Higgs boson below 200 GeV.

- Spectrum: $m_t = y_t \langle h \rangle$ $y_t = \frac{y_1 y_2}{\sqrt{y_1^2 + y_2^2}}$ $m_{t'} = \sqrt{y_1^2 + y_2^2} f$

$$t_{L,m} \approx t_L \quad t_{R,m} \approx (y_2 t_R - y_1 \psi_R) / \sqrt{y_1^2 + y_2^2}$$

$$t'_L \approx \psi_L \quad t'_{R,m} \approx (y_1 t_R + y_2 \psi_R) / \sqrt{y_1^2 + y_2^2}$$

- Large mixing between the right-handed parts of t and t' quarks.
- Both Z and B' couple to t_R and $t'_{R,m}$ with order one couplings.

Electroweak Precision Test

- At tree level, only the experimentally unmeasured top quark couplings to W and Z bosons are changed.
- At one-loop level, the strongest constraint comes from the T parameter:

$$\alpha T = \frac{3y_t^2 y_1^2 m_t^2}{16\pi^2 y_2^2 m_{t'}^2} \left(\log \frac{m_{t'}}{m_t} - 1 + \frac{y_1^2}{2y_2^2} \right)$$

[From PDG, $\alpha T < 1.2 \times 10^{-3}$ at 95% confidence level for $m_h < 300$ GeV.]


- For $y_1/y_2 < 3/4$, there is no bound on the symmetry breaking scale f . Hence, f can be as low as 400 GeV (to have the cutoff Λ above 5 TeV).

Signatures of the Z_2 Broken Model

- Two new parameters: y_2 and f (y_1 is determined by y_2 and y_t).
- Predicts two new particles: B' and t' .

$$M_{B'} = 5\sqrt{2}g'f/3 \approx 0.8f \quad m_{t'} = \sqrt{y_1^2 + y_2^2}f \geq 2f$$

- For $f \geq 400$ GeV, $M_{B'} \geq 300$ GeV. This possible light neutral gauge boson only couples to **top quarks** (nonuniversal).
- B' can mainly be produced through loop diagrams at Hadron Colliders like:[working in progress with Rakhi Mahbubani]



- B' decays to two top quarks. Mainly look for $t\bar{t} + 1$ jet.

Z_2 Unbroken Model

- To have a cold dark matter candidate, we need to keep this Z_2 to be unbroken to have stable particles. [Low and Cheng, 2003]
- Introduce two more vector-like quarks:

	$SU(3)_c$	$SU(2)_w$	$U(1)_1$	$U(1)_2$
H	1	2	1/2	1/2
S	1	1	5/3	-5/3
q_{1L}	3	2	1/6	1/6
t_R	3	1	2/3	2/3
b_R	3	1	-1/3	-1/3
ψ_{1L}	3	1	7/3	-1
ψ_{1R}	3	1	7/3	-1
ψ_{2L}	3	1	-1	7/3
ψ_{2R}	3	1	-1	7/3
q_{2L}	3	2	1/6	1/6
q_R	3	2	1/6	1/6

Gauge anomalies are cancelled.

Z_2 invariant

$$\begin{aligned}\mathcal{L}_t &= \frac{y_1}{\sqrt{2}} (\bar{q}_{1L} \tilde{H} + \bar{\psi}_{1L} S) t_R + y_2 f \bar{\psi}_{1L} \psi_{1R} \\ &+ \frac{y_1}{\sqrt{2}} (\bar{q}_{2L} \tilde{H} + \bar{\psi}_{2L} S^\dagger) t_R + y_2 f \bar{\psi}_{2L} \psi_{2R} \\ &+ \frac{y_3}{\sqrt{2}} f (\bar{q}_{1L} - \bar{q}_{2L}) q'_R + h.c.\end{aligned}$$

Under the Z_2 transformation, we have

$$Z_2 : \quad q_{1L} \leftrightarrow q_{2L}, \quad \psi_{1L,R} \leftrightarrow \psi_{2L,R}, \quad q'_R \rightarrow -q'_R, \\ B_1 \leftrightarrow B_2, \quad S \leftrightarrow S^\dagger$$

and all other fields are invariant

Mass Spectrum

Z_2 is exact; all particles are Z_2 eigenstates.

- Z_2 even particles:

$$\begin{aligned}
 t: \quad t_{L,m} &\approx t_L & t_{R,m} &\approx \frac{y_2 t_R - y_1 (\psi_{1R} + \psi_{2R}) / \sqrt{2}}{\sqrt{y_1^2 + y_2^2}} & m_t &= \frac{y_1 y_2}{y_1^2 + y_2^2} \langle h \rangle \\
 t'_+ : \quad t'_{+L} &\approx \frac{\psi_{1L} + \psi_{2L}}{\sqrt{2}} & t'_{+R} &\approx \frac{y_1 t_R + y_2 (\psi_{1R} + \psi_{2R}) / \sqrt{2}}{\sqrt{y_1^2 + y_2^2}} & m_{t'_+} &\approx \sqrt{y_1^2 + y_2^2} f \geq 2f
 \end{aligned}$$

The Λ^2 contribution to the Higgs mass from t is cancelled by t'_+ .

All other standard model particles are also Z_2 even.

- Z_2 odd particles:

$$\begin{aligned}
 t'_- : \quad t'_{-L} &\approx \frac{\psi_{1L} - \psi_{2L}}{\sqrt{2}} & t'_{-R} &\approx \frac{\psi_{1R} - \psi_{2R}}{\sqrt{2}} & m_{t'_-} &= y_2 f \\
 q'_- : \quad q'_{-L} &\approx \frac{q_{1L} - q_{2L}}{\sqrt{2}} & q'_{-R} &\approx q'_R & m_{q'_-} &= y_3 f \\
 B' : & & & (B_1 - B_2) / 2 & M_{B'} &\approx 0.8f
 \end{aligned}$$

- For $y_2, y_3 \geq 1$, B' is the lightest Z_2 odd particle and a potential **dark matter candidate** in this model.

Dark Matter

- From WMAP, the relic abundance of the dark matter is:

$$0.098 < \Omega_{dm} h^2 < 0.122 (2\sigma)$$


- In the non-relativistic limit, $\Omega_{dm} h^2$ is relating to sum of the quantities, $a(X) = v_r \sigma(B' B' \rightarrow X)$, as

$$\Omega_{dm} h^2 \approx \frac{1.04 \times 10^9 \text{ GeV}^{-1}}{M_{pl}} \frac{x_F}{\sqrt{g^*}} \frac{1}{a_{tot}}$$

- Approximately, only need to calculate a_{tot} and require:

$$a_{tot} \approx 0.81 \pm 0.09 \text{ pb}$$

Couplings of B' to Higgs Boson



Minimal Little Higgs
Model


$$\frac{50}{9} g'^2 v$$

Hypercharge-like Gauge
Boson [LHT and UED]


$$\frac{1}{2} g'^2 v$$

Relic Abundance

The leading processes for $B' B'$ annihilation into SM particles:




$$a(WW) = 2a(ZZ) = 2a(h^0 h^0) = \frac{2\pi\alpha^2}{3\cos^4\theta_w} \frac{5^4}{3^4} \frac{1}{M_{B'}^2}$$



$$a(tt) = \frac{16\pi\alpha^2}{3\cos^4\theta_w} \frac{5^4}{3^4} \frac{y_t^4}{y_2^4} \frac{M_{B'}^2}{(M_{B'}^2 + m_{t'}^2)^2}$$

For $y_2 \gg 1$, $a(\bar{t}t)$ is negligible.




$$[M_{B'} \approx 0.8 f m_{t'} = y_2 f]$$

$$0.098 < \Omega_{dm} h^2 < 0.122 (2\sigma) \Rightarrow a_{tot} \approx 0.81 \pm 0.09 \text{ pb}$$

Direct Detection

- Measure the recoil energy in the elastic scattering of dark matter particles with nuclei.





Only contribute to spin-independent cross section

- Using the matrix element of quarks and gluons in a nucleon state: [Ellis, Olive, Santos, Spanos, 2005]

$$\sigma_{SI} \approx \frac{0.35^2 g'^4}{16\pi M_{B'}^2} \frac{10^4}{3^4} \frac{m_p^4}{m_h^4} \approx 1.6 \times 10^{-44} \text{cm}^2 \left(\frac{1 \text{ TeV}}{M_{B'}}\right)^2 \left(\frac{100 \text{ GeV}}{m_h}\right)^4$$

- Box diagrams with the top quark propagating in the the loop also contribute to spin-dependent cross section.

Direct Detection



Summary

- A very simple little Higgs model has been constructed based on the $SU(2)_w \times U(1)^2$ gauge symmetry.
- A Z_2 interchanging symmetry is introduced between these two $U(1)$'s.
- For Z_2 **broken** case: only B' and t' appears in the EFT. The mass of B' can be as light as 300 GeV.
- For Z_2 **unbroken** case:
 - B' is a stable particle and can serve as a dark matter candidate.
 - The direct detection of this B' dark matter is promising.
 - The $\sigma_{SI}(B'N)$ is two order of magnitude larger than a hypercharge-like neutral gauge boson dark matter candidate.