## Astroparticle Physics in the LHC Era





#### John Beacom The Ohio State University



John Beacom, The Ohio State University

#### Plan of the Talk

Cosmic Rays, Gamma Rays, and Neutrinos Gamma Ray Detectors and Sources Neutrino Detectors and Sources Prospecting for New Physics Concluding Perspectives

#### Cosmic Rays, Gamma Rays, and Neutrinos

## Are there high energy processes in nature? Do these produce gamma rays and neutrinos?

### What's So Special About a TeV?


1 TeV = 10<sup>12</sup> eV = 1.6 erg per particle Far above atomic (eV) and nuclear (MeV) scales



John Beacom, The Ohio State University

Pheno Symposium, University of Wisconsin-Madison, April 2008

#### Cosmic Ray Protons, Electrons, and Nuclei



#### Cosmic Rays Imply Gamma Rays and Neutrinos

Hadronic mechanism

$$p + p \rightarrow p + p + \pi^{0}, p + n + \pi^{1}$$
$$\pi^{0} \rightarrow 2\gamma, \pi^{\pm} \rightarrow e^{\pm} + 3v$$

Leptonic mechanism

$$e^- + \gamma \rightarrow \gamma + e^-$$


• Nuclear (A\*) mechanism

$$A' + \gamma \to A^* + X$$
$$A^* \to A + \gamma$$

Anchordoqui, Beacom, Goldberg, Palomares-Ruiz, Weiler, PRL 98, 121101 (2007)

## Astronomy with New Messengers

cosmic rays	gamma rays	neutrinos
energetic	direct	revealing
divertable	stoppable	untrustworthy?



John Beacom, The Ohio State University

Pheno Symposium, University of Wisconsin-Madison, April 2008

## Gamma Ray Detectors and Sources


Do luminous high energy gamma ray sources exist? Can we find them and measure them?

#### **Gamma-Ray Detection Techniques**


below ~ 0.3 TeV

~ 0.3-30 TeV


above ~ 3 TeV



primary gamma ray




air Cerenkov from shower




# shower at ground

#### **EGRET Source Results**



#### **EGRET Diffuse Results**



#### **HESS Observatory**

#### Four 13-m telescopes operated synchronously




#### In full operation in Namibia since 2004


John Beacom, The Ohio State University

#### **Resolved Extended Sources**

#### supernova remnant RX J1713.7-3946



#### HESS Survey of the Inner Galaxy



#### **VERITAS** Observatory




#### Instrument:

- Four 12-m telescopes
- 500-pixel cameras (3.5° FoV)
- FLWO, Mt. Hopkins, AZ (1268 m)
- Completed Spring 2007

#### Specifications:

- Energy threshold
- Source location
- Energy resolution
- ~ 150 GeV
- < 0.05°
  - ~ 10-20 %


#### Skymap of VHE Gamma-Ray Sources



2008-03-16 - Up-to-date plot available at http://www.mppmu.mpg.de/~rwagner/sources/

John Beacom, The Ohio State University


#### Milagro Experiment



- Water Cherenkov Detector
- 2600m asl
- 898 detectors
  - 450(t)/273(b) in pond
  - 175 water tanks
- $\sim$  4000 m<sup>2</sup> / 4.0x10<sup>4</sup> m<sup>2</sup>
- 2-20 TeV median energy
- 1700 Hz trigger rate
- 0.4°-1.0° resolution
- 95% background rejection


#### Milagro 12 TeV Diffuse

#### First partial preview of the Northern neutrino sky



Milagro Collaboration (2007)


#### Cygnus Region



#### Beacom, Kistler (2007)

#### Milagro J2019+37 Region

#### MGRO J2019+37 in Cygnus



#### Gamma-Ray Scorecard

Up to ~ 0.1 TeV
EGRET saw ~ 10<sup>2</sup> sources in the full sky
EGRET saw full-sky diffuse emission

- Around 1 TeV Whipple, HESS, etc saw tens of sources No data on diffuse emission
- Around 10 TeV

Milagro saw a few sources *in survey mode* Milagro saw diffuse emission in part of sky

#### Neutrino Detectors and Sources

# Do luminous high energy neutrino sources exist? Can we find them and measure them?

#### **Muon-Induced Neutrinos**


• For hadronic sources,  $\phi_{v} \sim \phi_{v}$ 


• Detection reaction is  $v_{\mu} + n \rightarrow \mu^- + p$ 

Muon range is ~ 1-10 km

• Near 1 TeV,  $P(v \rightarrow \mu) \sim n\sigma L \sim 10^{-6}$ 








John Beacom, The Ohio State University

Pheno Symposium, University of Wisconsin-Madison, April 2008


## Neutrino Skymap?



#### AMANDA Collaboration (2003)

John Beacom, The Ohio State University


#### **AMANDA Results**



#### AMANDA Collaboration (2007)

John Beacom, The Ohio State University

#### **Probing Sources With Neutrinos**



Kistler, Beacom (2006)

Definitive sign of hadronic mechanism


km<sup>3</sup> detectors are big enough

Advantages at large energies

Neutrino-only sources?

#### Galactic Neutrino Sources

#### Vela Jr. supernova remnant




#### (and many more)

Source	$\phi_{\gamma}$	Γ	$E_{v}^{cut}$ (TeV)	$N_{\mu}(> 1 \text{ TeV})$
Vela Jr.	21.0	2.1	10	3.1
(RX J0852.0-4622)			25	4.9
			50	6.1
GC Diffuse	5.2	2.29	20	0.5
			50	0.7
(+ GC Source)			20	0.8
			50	1.0
RX J1713.7–3946	15.0	2.19	50	2.8
	20.4	1.98	6	2.2
Vela X	9.0	1.45	7	4.5
Crab (IceCube)	33.0	2.57	50	2.7
HESS J1514–591	5.7	2.27	25	0.9
			50	1.1
HESS J1616-508	6.0	2.35	10	0.5
			50	0.9
HESS J1632-478	5.5	2.12	10	0.8
			50	1.5

#### Kistler, Beacom (2006)

#### Neutrinos from the Milagro Source

#### MGRO J2019+37 in Cygnus



Beacom, Kistler (2007)

#### Neutrino Scorecard

 Up to ~ 1 TeV
Super-Kamiokande, other experiments saw only atmospheric neutrinos

- Above 1 TeV AMANDA saw only atmospheric neutrinos Excellent prospects for IceCube
- At much higher energies From several experiments, only upper limits on fluxes

Prospecting for New Physics

Do neutrinos or dark matter have new properties?


Are there dark matter annihilation signals?

What surprises are out there?

# Neutrino Flavor RatiosMocioiu $\pi \rightarrow \mu + v_{\mu} \rightarrow e + v_{e} + 2v_{\mu}$ $\phi_{e} : \phi_{\mu} : \phi_{\tau} \sim 1 : 2 : 0$ $v_{\mu} \leftrightarrow v_{\tau}$ $\phi_{e} : \phi_{\mu} : \phi_{\tau} \sim 1 : 1 : 1$

Neutrino invisible decays are not ruled out, and would greatly alter the ratios

Other new physics can lead to different ratios



Beacom, Bell, Hooper, Pakvasa, Weiler, PRL 90, 181301 (2003); Beacom, Bell, Hooper, Pakvasa, Weiler, PRD 69, 017303 (2004)


John Beacom, The Ohio State University

#### Dark Matter Properties Profumo

Albuquerque, Burdman, Chacko, PRL 92, 221802 (2004)

NLSP is charged and short-lived

Astro neutrinos make NSLP pairs in Earth




Energetic NLSP pairs make a new signal in IceCube Better if prompt atmospheric neutrino flux is large Ando, Beacom, Profumo, Rainwater, JCAP04 (2008) 029

John Beacom, The Ohio State University

## Profumo Dark Matter Annihilation Dent, Mack

#### Annihilation products: gamma rays and neutrinos



## **Concluding Perspectives**


#### Conclusions

Luminous TeV gamma-ray sources exist: Most have uncertain astronomical associations Fundamental question of production mechanism

Better gamma-ray observations are essential: Increase energy range to test spectra Refine angular resolution to make identifications Cover the full sky to study populations and diffuse

Neutrino observations can be decisive: IceCube and other detectors coming online A novel probe of the cosmos and new particle physics

Great future with lots of complementary data



# Divide by zero?

Stand by ... It's about to happen.

John Beacom, The Ohio State University